

# (12) UK Patent Application (19) GB (11) 2 282 598 (13) A

(43) Date of A Publication 12.04.1995

(21) Application No 9420114.2

(22) Date of Filing 05.10.1994

(30) Priority Data

(31) 133493

(32) 08.10.1993

(33) US

(71) Applicant(s)

Merck & Co Inc

(Incorporated in USA - New Jersey)

P O Box 2000, 126 East Lincoln Avenue, Rahway,  
New Jersey 07065-0900, United States of America

(72) Inventor(s)

William K Hagmann

Charles G. Caldwell

Paul R Gooley

(74) Agent and/or Address for Service

W G Cole

Merck & Co Inc, European Patent Department,  
Terlings Park, Eastwick Road, HARLOW, Essex,  
CM20 2QR, United Kingdom

(51) INT CL<sup>6</sup>

C07C 235/82, A61K 31/19 31/215

(52) UK CL (Edition N)

C2C CAA CKZ C1175 C1176 C200 C22X C22Y C221  
C225 C226 C227 C280 C282 C30Y C304 C34Y C342  
C366 C367 C368 C575 C699 C80Y C802  
U1S S1313 S2412 S2415 S2416

(56) Documents Cited


None

(58) Field of Search

UK CL (Edition M) C2C CKK CKN CKS CKZ  
INT CL<sup>5</sup> C07C, C07D  
ONLINE DATABASE: CAS

## (54) Substituted cyclic derivatives as novel antidegenerative agents

(57) Novel substituted cyclic compounds of Formula I are found to be useful inhibitors of matrix metalloendoproteinase-mediated diseases including osteoarthritis, rheumatoid arthritis, septic arthritis, tumor invasion in certain cancers, periodontal disease, corneal ulceration, proteinuria, dystrophic epidermolysis bullosa, and coronary thrombosis associated with atherosclerotic plaque rupture. The matrix metalloendoproteinases are a family of zinc-containing proteinases including but not limited to stromelysin, collagenase, and gelatinase, that are capable of degrading the major components of articular cartilage and basement membranes. The inhibitors claimed herein may also be useful in preventing the pathological sequelae following a traumatic injury that could lead to a permanent disability. These compounds may also be useful as novel birth control agents by preventing ovulation or implantation.



I


GB 2 282 598 A

TITLE OF THE INVENTIONSUBSTITUTED CYCLIC DERIVATIVES AS NOVEL  
ANTIDEGENERATIVE AGENTS5 BACKGROUND OF THE INVENTION1. Field of the Invention

By this invention there is provided novel substituted  
cyclic compounds of Formula I which are useful inhibitors of  
10 matrix metalloendoproteinase-mediated diseases including  
osteoarthritis, rheumatoid arthritis, septic arthritis, tumor invasion  
in certain cancers, periodontal disease, corneal ulceration,  
proteinuria, dystrophic epidermolysis bullosa, and coronary  
thrombosis associated with atherosclerotic plaque rupture. The  
15 matrix metalloendoproteinases are a family of zinc-containing  
proteinases including but not limited to stromelysin, collagenase, and  
gelatinase, that are capable of degrading the major components of  
articular cartilage and basement membranes. The inhibitors claimed  
herein may also be useful in preventing the pathological sequelae  
20 following a traumatic injury that could lead to a permanent  
disability. These compounds may also be useful as novel birth  
control agents by preventing ovulation or implantation.

25

30



I

## 2. Brief Description of Disclosures in the Art

The disability observed in osteoarthritis (OA) and rheumatoid arthritis (RA) is largely due to the loss of articular cartilage. No therapeutic agent in the prior art is known to prevent the attrition of articular cartilage in these diseases.

"Disease modifying antirheumatic drugs" (DMARD), i.e., agents capable of preventing or slowing the ultimate loss of joint function in OA and RA are widely sought. Generic nonsteroidal antiinflammatory drugs (NSAIDs) may be combined with such agents to provide some relief from pain and swelling.

Stromelysin (aka. proteoglycanase, matrix metalloproteinase-3, MMP-3, procollagenase activator, "transin"), collagenase (aka. interstitial collagenase, matrix metalloproteinase-1), and gelatinase (aka. type IV collagenase, matrix metalloproteinase-2, MMP-2, 72kDa-gelatinase, gelatinase A or type V collagenase, matrix metalloproteinase-9, MMP-9, 95kDa-gelatinase, gelatinase B) are metalloendoproteinases secreted by fibroblasts, chondrocytes, and macrophage and are capable of degrading the major connective tissue components of articular cartilage or basement membranes. Elevated levels of these enzymes have been detected in joints of arthritic humans and animals: K.A. Hastay, R.A. Reife, A.H. Kang, J.M. Stuart, "The role of stromelysin in the cartilage destruction that accompanies inflammatory arthritis", *Arthr. Rheum.*, **33**, 388-97 (1990); S.M. Krane, E.P. Amento, M.B. Goldring, S.R. Goldring, and M.L. Stephenson, "Modulation of matrix synthesis and degradation in joint inflammation", in "The Control of Tissue Damage", A.B. Glauert (ed.), Elsevier Sci. Publ., Amsterdam, 1988, Ch. 14, pp 179-95; A. Blanckaert, B. Mazieres, Y. Eeckhout, G. Vaes, "Direct extraction and assay of collagenase from human osteoarthritic cartilage", *Clin. Chim. Acta*, **185** 73-80 (1989); J.-P. Pelletier, M.-P. Faure, J. A. DiBattista, S. Wilhelm, D. Visco, J. Martel-Pelletier, "Coordinate Synthesis of Stromelysin, Interleukin-1, and Oncogene Proteins in Experimental Osteoarthritis", *Am. J. Path.* **142**, 95-105 (1993); L. A. Walakovits,

N. Bhardwaj, G. S. Gallick, M. W. Lark, "Detection of high levels of stromelysin and collagenase in synovial fluid from patients with rheumatoid arthritis and post-traumatic knee injury", *Arthr. Rheum.* **35**, 35-42 (1992). Each enzyme is secreted from these cells as an inactive proenzyme which is subsequently activated. There is evidence that stromelysin may be the in vivo activator for collagenase and gelatinase, implying a cascade for degradative enzyme activity: A. Ho, H. Nagase, "Evidence that human rheumatoid synovial matrix metalloproteinase 3 is an endogenous activator of procollagenase", *Arch Biochem Biophys.*, **267**, 211-16 (1988); G. Murphy, M.I. Crockett, P.E. Stephens, B.J. Smith, A.J.P. Docherty, "Stromelysin is an activator of procollagenase", *Biochem. J.*, **248**, 265-8 (1987); Y. Ogata, J. J. Enghild, H. Nagase, "Matrix Metalloproteinase 3 (stromelysin) activates the precursor for the human matrix metalloproteinase 9", *J. Biol. Chem.* **267**, 3581-3584 (1992); K. Mikazaki, F. Umenishi, K. Funahashi, N. Yasumitsu, M. Umeda, "Activation of TIMP-2/Progelatinase A Complex by Stromelysin", *Biochem. Biophys. Res. Commun.*, **185**, 852-859 (1992). Inhibiting stromelysin could limit the activation of collagenase and gelatinase as well as prevent the degradation of proteoglycan.

That stromelysin inhibition may be effective in preventing articular cartilage degradation has been demonstrated in vitro by measuring the effect of matrix metalloendoproteinase inhibitors on proteoglycan release from rabbit cartilage explants: C.B. Caputo, L.A. Sygowski, S.P. Patton, D.J. Wolanin, A. Shaw, R.A. Roberts, G. DiPasquale, *J. Orthopaedic Res.*, **6**, 103-8 (1988).

There is an extensive literature on the involvement of these metalloproteinases in arthritis, but there is very little to guide one in developing a specific inhibitor for each enzyme.

In preliminary studies of rabbit proteoglycanase with substrates and inhibitors, little was found to indicate the enzyme's requirements for hydrolysis or inhibition beyond a preference for hydrophobic residues at the P<sub>1</sub>' position: A. Shaw, R.A. Roberts,

D.J. Wolanin, "Small substrates and inhibitors of the metalloproteoglycanase of rabbit articular chondrocytes", *Adv. Inflam. Res.*, **12**, 67-79 (1988).

Human rheumatoid synovial collagenase has been shown  
5 to share ~ 50% homology with human stromelysin: S.E. Whitham, G. Murphy, P. Angel, H.J. Rahmsdorf, B.J. Smith, A. Lyons, T.J.R. Harris, J.J. Reynolds, P. Herrlich, A.J.P. Docherty, "Comparison of human stromelysin and collagenase by cloning and sequence  
10 analysis", *Biochem. J.*, **240**, 913-6 (1986). Many collagenase inhibitors have been designed around the cleavage site of the  $\alpha$ -chain sequence of Type II collagen: W.H. Johnson, N.A. Roberts, N. Brokakoti, "Collagenase inhibitors: their design and potential  
15 therapeutic use", *J. Enzyme Inhib.*, **2**, 1-22 (1987); M. A. Schwartz, H. E. Van Wart, "Synthetic Inhibitors of Bacterial and Mammalian Interstitial Collagenases", in *Progress in Medicinal Chemistry*, vol  
20 29, G. P. Ellis, D. K. Luscombe (eds.), Elsevier Sci. Publish., Amsterdam, 1992, pp 271-334. One such inhibitor, N-[3-(benzyloxycarbonyl)amino-1-carboxy-n-propyl]-L-leucyl-O-methyl-L-tyrosine, N-methylamide, prepared at G.D. Searle, Inc., and  
25 shown to be a potent inhibitor of human rheumatoid synovial collagenase ( $IC_{50} = 0.8 \mu M$ ), was also found to inhibit rabbit bone proteoglycanase ( $IC_{50} = 0.5 \mu M$ ): J.-M. Delaisse, Y. Eeckhout, C.Sear, A. Galloway, K. McCullagh, G. Vaes, "A new synthetic inhibitor of mammalian tissue collagenase inhibits bone resorption in  
culture", *Biochem. Biophys. Res. Commun.*, **133**, 483-90 (1985).

Gelatinase-A (MR ~ 72,000) has been isolated from  
rheumatoid fibroblasts: Y. Okada, T. Morodomi, J. J. Enghild, K. Suzuki, A. Yasui, I. Nakanishi, G. Salvesen, H. Nagase, "Matrix  
30 metalloproteinase 2 from human rheumatoid synovial fibroblasts", *Eur. J., Biochem.*, **194**, 721-30 (1990). The synthesis of the proenzyme of this matrix metalloproteinase is not coordinately regulated with stromelysin or collagenase and its activation may also be different. The role of gelatinase-A in the tissue destruction of articular cartilage appears different from the other two enzymes and,

therefore, its inhibition may provide additional protection from degradation. A higher molecular weight gelatinase-B (MR ~95,000; aka. type-V collagenase, matrix metalloproteinase-9, MMP-9) is also secreted by fibroblasts and monocytes and may be involved in cartilage degradation.

As appreciated by those of skill in the art, the significant proportion of homology between human fibroblast collagenase, stromelysin, and gelatinase leads to the possibility that a compound that inhibits one enzyme may to some degree inhibit all of them.

The applicants believe that matrix metalloproteinase inhibitors have utility in preventing articular cartilage damage associated with septic arthritis. Bacterial infections of the joints can elicit an inflammatory response that may then be perpetuated beyond what is needed for removal of the infective agent resulting in permanent damage to structural components. Bacterial agents have been used in animal models to elicit an arthritic response with the appearance of proteolytic activities. See J.P. Case, J. Sano, R. Lafyatis, E.F. Remmers, G.K. Kumkumian, R.L. Wilder, "Transin/stromelysin expression in the synovium of rats with experimental erosive arthritis", *J. Clin Invest.*, **84**, 1731-40 (1989); R.J. Williams, R.L. Smith, D.J. Schurman, "Septic Arthritis: Staphylococcal induction of chondrocyte proteolytic activity", *Arthr. Rheum.*, **33**, 533-41 (1990).

The applicants also believe that inhibitors of stromelysin, collagenase, and gelatinase will be useful to control tumor metastasis, optionally in combination with current chemotherapy and/or radiation. See L.M. Matrisian, G.T. Bowden, P. Krieg, G. Furstenberger, J.P. Briand, P. Leroy, R. Breathnach, "The mRNA coding for the secreted protease transin is expressed more abundantly in malignant than in benign tumors", *Proc. Natl. Acad. Sci., USA*, **83**, 9413-7 (1986); S.M. Wilhelm, I.E. Collier, A. Kronberger, A.Z. Eisen, B.L. Marmer, G.A. Grant, E.A. Bauer, G. I. Goldberg, "Human skin fibroblast stromelysin: structure, glycosylation, substrate specificity, and differential expression in

normal and tumorigenic cells", *Ibid.*, **84**, 6725-29 (1987); Z. Werb et al., Signal transduction through the fibronectin receptor induces collagenase and stromelysin gene expression, *J. Cell Biol.*, **109**, 872-889 (1989); L.A. Liotta, C.N. Rao, S.H. Barsky, "Tumor invasion and the extracellular matrix", *Lab. Invest.*, **49**, 636-649 (1983); R. Reich, B. Stratford, K. Klein, G. R. Martin, R. A. Mueller, G. C. Fuller, "Inhibitors of collagenase IV and cell adhesion reduce the invasive activity of malignant tumor cells", in "Metastasis: Ciba Foundation Symposium"; Wiley, Chichester, 1988, pp. 193-210.

Secreted proteinases such as stromelysin, collagenase, and gelatinase play an important role in processes involved in the movement of cells during metastatic tumor invasion. Indeed, there is also evidence that the matrix metalloproteinases are overexpressed in certain metastatic tumor cell lines. In this context, the enzyme functions to penetrate underlying basement membranes and allow the tumor cell to escape from the site of primary tumor formation and enter circulation. After adhering to blood vessel walls, the tumor cells use these same metalloendoproteinases to pierce underlying basement membranes and penetrate other tissues, thereby leading to tumor metastasis. Inhibition of this process would prevent metastasis and improve the efficacy of current treatments with chemotherapeutics and/or radiation.

These inhibitors should also be useful for controlling periodontal diseases, such as gingivitis. Both collagenase and stromelysin activities have been isolated from fibroblasts isolated from inflamed gingiva: V.J. Uitto, R. Applegren, P.J. Robinson, "Collagenase and neutral metalloproteinase activity in extracts of inflamed human gingiva", *J. Periodontal Res.*, **16**, 417-424 (1981). Enzyme levels have been correlated to the severity of gum disease: C.M. Overall, O.W. Wiebkin, J.C. Thonard, "Demonstration of tissue collagenase activity in vivo and its relationship to inflammation severity in human gingiva", *J. Periodontal Res.*, **22**, 81-88 (1987).

Proteolytic processes have been observed in the ulceration of the cornea following alkali burns: S.I. Brown, C.A.

Weller, H.E. Wasserman, "Collagenolytic activity of alkali-burned corneas", *Arch. Ophthalmol.*, **81**, 370-373 (1969). Mercapto-containing peptides inhibit the collagenase isolated from alkali-burned rabbit cornea: F.R. Burns, M.S. Stack, R.D. Gray, C.A. Paterson, *Invest. Ophthalmol.*, **30**, 1569-1575 (1989). Treatment of alkali-burned eyes or eyes exhibiting corneal ulceration as a result of infection with inhibitors of these metalloendoproteinases in combination with sodium citrate or sodium ascorbate and/or antimicrobials may be effective in preventing developing corneal degradation.

Stromelysin has been implicated in the degradation of structural components of the glomerular basement membrane (GBM) of the kidney, the major function of which is to restrict passage of plasma proteins into the urine; W.H. Baricos, G. Murphy, Y. Zhou, H.H. Nguyen, S.V. Shah, "Degradation of glomerular basement membrane by purified mammalian metalloproteinases", *Biochem. J.*, **254**, 609-612 (1988). Proteinuria, a result of glomerular disease, is excess protein in the urine caused by increased permeability of the GBM to plasma proteins. The underlying causes of this increased GBM permeability are unknown, but proteinases including stromelysin may play an important role in glomerular diseases. Inhibition of this enzyme may alleviate the proteinuria associated with kidney malfunction.

It is suggested that inhibition of stromelysin activity may prevent the rupturing of atherosclerotic plaques leading to coronary thrombosis. The tearing or rupture of atherosclerotic plaques is the most common event initiating coronary thrombosis. Destabilization and degradation of the connective tissue matrix surrounding these plaques by proteolytic enzymes or cytokines released by infiltrating inflammatory cells has been proposed as a cause of plaque fissuring. Such tearing of these plaques can cause an acute thrombotic event as blood rapidly flows out of the blood vessel. High levels of stromelysin RNA message have been found to be localized to individual cells in atherosclerotic plaques removed from heart



transplant patients at the time of surgery: A. M. Henney, P. R. Wakeley, M. J. Davies, K. Foster, R. Hembry, G. Murphy, S. Humphries, "Localization of stromelysin gene expression in atherosclerotic plaques by *in situ* hybridization", *Proc. Nat'l. Acad. Sci. USA*, **88**, 8154-8158 (1991). Inhibition of stromelysin by these compounds may aid in preventing or delaying the degradation of the connective tissue matrix that stabilizes the atherosclerotic plaques, thereby preventing events leading to acute coronary thrombosis.

Collagenolytic and stromelysin activity have also been observed in dystrophic epidermolysis bullosa: A. Kronberger, K.J. Valle, A.Z. Eisen, E.A. Bauer, *J. Invest. Dermatol.*, **79** 208-211 (1982); D. Sawamura, T. Sugawara, I. Hashimoto, L. Bruckmer-Tuderman, D. Fujimoto, Y. Okada, N. Utsumi, H. Shikata, *Biochem. Biophys. Res. Commun.*, **174**, 1003-8 (1991). Inhibition of metalloendoproteinases should limit the rapid destruction of connective components of the skin.

It is also believed that specific inhibitors of stromelysin and collagenase should be useful as birth control agents. There is evidence that expression of metalloendoproteinases, including stromelysin and collagenase, is observed in unfertilized eggs and zygotes and at further cleavage stages and increased at the blastocyst stage of fetal development and with endoderm differentiation: C.A. Brenner, R.R. Adler, D.A. Rappolee, R.A. Pedersen, Z. Werb, "Genes for extracellular matrix-degrading metalloproteinases and their inhibitor, TIMP, are expressed during early mammalian development", *Genes & Develop.*, **3**, 848-59 (1989). By analogy to tumor invasion, a blastocyst may express metalloproteinases in order to penetrate the extracellular matrix of the uterine wall during implantation. Inhibition of stromelysin and collagenase during these early developmental processes should presumably prevent normal embryonic development and/or implantation in the uterus. Such intervention would constitute a novel method of birth control. In addition there is evidence that collagenase is important in ovulation processes. In this example, a covering of collagen over the apical

region of the follicle must be penetrated in order for the ovum to escape. Collagenase has been detected during this process and an inhibitor has been shown to be effective in preventing ovulation: J.F. Woessner, N. Morioka, C. Zhu, T. Mukaida, T. Butler, W.J. LeMaire "Connective tissue breakdown in ovulation", *Steroids*, **54**, 491-499 (1989). There may also be a role for stromelysin activity during ovulation: C.K.L. Too, G.D. Bryant-Greenwood, F.C. Greenwood, "Relaxin increases the release of plasminogen activator, collagenase, and proteo-glycanase from rat granulosa cells in vitro", *Endocrin.*, **115**, 1043-1050 (1984).


Metalloendoproteinases have also been implicated in the cleavage of proteins associated with the myelin sheath surrounding nerve fibers: A. Chantry, C. Earl, N. Groome, P. Glynn, "Metalloendoprotease Cleavage of 18.2- and 14.1-kilodalton basic proteins dissociating from rodent myelin membranes generates 10.0- and 5.9kilodalton C-terminal fragments", *J. Neurochem.* **50**, 688-694 (1988). Inhibitors of such processes may prevent nerve sheath degeneration associated with several neurological diseases, including muscular dystrophy.

In addition to extracellular matrix comprising structural components, stromelysin can degrade other in vivo substrates including the inhibitors  $\alpha_1$ -proteinase inhibitor and may therefore influence the activities of other proteinases such as elastase: P. G. Winyard, Z. Zhang, K. Chidwick, D. R. Blake, R. W. Carrell, G. Murphy, "Proteolytic inactivation of human  $\alpha_1$ -antitrypsin by human stromelysin", *FEBS Letts.*, **279**, 1, 91-94 (1991). Inhibition of the matrix metalloendoproteinases may potentiate the antiproteinase activity of these endogenous inhibitors.

### SUMMARY OF THE INVENTION

By this invention there is provided a class of novel substituted cyclic compounds of the following Formula I which are useful inhibitors of matrix metalloendoproteinase-mediated diseases

including degenerative diseases (such as defined above) and certain cancers:



15 wherein:

n is 1 - 4;

p is 1-2;

20

Ar is an aryl or heteroaryl group selected from the group consisting of:

- (1) phenyl,
- (2) naphthyl,
- 25 (3) pyridyl,
- (4) pyrrol,
- (5) furyl,
- (6) thienyl,
- (7) isothiazolyl,
- 30 (8) imidazolyl,
- (9) benzimidazolyl,
- (10) tetrazolyl,
- (11) pyrazinyl,
- (12) pyrimidyl,
- (13) quinolyl,

- 5 (14) isoquinolyl,  
(15) benzofuryl,  
(16) isobenzofuryl,  
(17) benzothienyl,  
(18) pyrazolyl,  
(19) indolyl,  
(20) isoindolyl,  
(21) purinyl,  
10 (22) carboxazolyl,  
(23) isoxazolyl,  
(24) thiazolyl,  
(25) oxazolyl,  
(26) benzthiazolyl, and  
(27) benzoxazolyl,

15 which can be mono- or di-substituted with substituents  
independently selected from C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, halo,  
hydroxy, amino, C<sub>1-6</sub>alkylamino, aminoC<sub>1-6</sub>alkyl, carboxyl,  
carboxylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl, aryl or heteroaryl as defined  
20 above;


R<sub>1</sub> is NHOH, OH, OC<sub>1-10</sub>alkyl, OCH(C<sub>1-10</sub>alkyl)OC(O)C<sub>1-10</sub>alkyl,  
OCH<sub>2</sub>phenyl, wherein the phenyl group may be substituted  
with hydrogen, carboxy, carboxyC<sub>1-3</sub>alkyl, aminocarbonyl,  
C<sub>1-6</sub> alkylaminocarbonyl;

25


R<sub>2</sub> is hydrogen or a mono- or di-substituted C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy,  
or C<sub>2-6</sub>alkenyl wherein the substituents are independently  
selected from the group consisting of:

- 30 (a) hydrogen,  
(b) carboxy,  
(c) aminocarbonyl,  
(d) C<sub>1-6</sub>alkoxy,  
(e) C<sub>1-6</sub>alkylcarbonyl,  
(f) aryl or heteroaryl, as defined above;

- (g) aryloxy or heteroaryloxy wherein the aryl or heteroaryl groups are defined above;
- (h) aroyl or heteroaroyl wherein the aryl or heteroaryl groups are defined above;




- (i) wherein  $R_a$  and  $R_b$  are each independently hydrogen; aryl or heteroaryl and mono and di-substituted aryl or heteroaryl as defined above; or substituted  $C_{1-6}$ alkyl wherein the substituent is selected from hydroxy, halo, and aryl or heteroaryl, or wherein  $R_a$  and  $R_b$  are joined together with the nitrogen and carbon atoms to which they are attached, to form a lactam or benzolactam ring wherein the lactam portion thereof is a ring of 5, 6, 7, or 8 atoms, said lactam or benzolactam containing a single hetero atom;



- (j) wherein  $R_a$  and  $R_b$  are each independently hydrogen; aryl or heteroaryl and mono and di-substituted aryl or heteroaryl as defined above ; or substituted  $C_{1-6}$ alkyl wherein the substituent is selected from hydroxy, halo, and aryl, or wherein  $R_a$  and  $R_b$  are joined together with the nitrogen and carbon atoms to which they are attached to form a cyclic imide, or is benzofused onto a cyclic imide, wherein the cyclic imide portion thereof is a ring of 5, 6, 7, or 8 atoms, said cyclic imide containing a single hetero atom;

- (k) amino or mono- or di-substituted amino wherein the substituents are independently selected from C1-6 alkyl and aryl or heteroaryl as defined above;


5



- (l) wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above ;  
or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen and oxygen atoms to which they are attached, to form a cyclic-carbamate, or a benzofused cyclic-carbamate, wherein the carbamate ring contains 5, 6, 7, or 8 atoms, said ring containing the two heteroatoms;

10


15



- (m) wherein R<sub>a</sub>, R<sub>b</sub>, and R<sub>c</sub> are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen atoms to which they are attached to form a cyclic-urea or benzofused cyclic-urea, said urea ring containing 5, 6, 7, or 8 atoms and said two nitrogens;


20

25




- (n) wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen and sulfur atoms to which they are attached, to form a cyclic-sulfonamide or benzofused cyclic-sulfonamide ring, said sulfonamide ring containing 5, 6, 7, or 8 atoms and said heteroatoms;

30



- (o) wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms and said nitrogen atom;



- (p) wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms and said nitrogen atom;

$\text{R}_3$  is

- (a) mono- or di-substituted C1-10alkylamino, or mono- or di-substituted di-C1-10alkyl amino wherein the substituents are selected from the group consisting of:
- (1) hydrogen,
  - (2) mercapto,
  - (3) hydroxy,
  - (4) carboxy,
  - (5) amino,
  - (6) aminocarbonyl,
  - (7) mono- or di-C1-6alkyl amino,
  - (8) mono- or di-C1-6alkyl aminocarbonyl,
  - (9) guanidino,
  - (10) aryl or heteroaryl, as defined above;


(b) an amino acid derivative of Formula II



wherein Rf and Rg are individually selected from:

- 10
- (a) hydrogen,
  - (b) C<sub>1</sub>-6alkyl,
  - (c) mercapto C<sub>1</sub>-6alkyl,
  - (d) hydroxy C<sub>1</sub>-6alkyl,
  - (e) carboxy C<sub>1</sub>-6alkyl,
  - 15 (f) amino C<sub>1</sub>-6alkyl
  - (g) aminocarbonyl C<sub>1</sub>-6alkyl,
  - (h) mono- or di-C<sub>1</sub>-6alkyl amino C<sub>1</sub>-6alkyl,
  - (i) guanidino C<sub>1</sub>-6alkyl,
  - (j) substituted phenyl C<sub>1</sub>-6alkyl, wherein the substituent is
  - 20 hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
  - (k) substituted indolyl C<sub>1</sub>-6alkyl, wherein the substituent is
  - hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
  - (l) substituted imidazolyl C<sub>2</sub>-6alkyl wherein the substituent is
  - hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
  - 25 (m) substituted pyridyl C<sub>1</sub>-6alkyl wherein the substituent is
  - hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
  - (n) substituted pyridylamino C<sub>1</sub>-6alkyl wherein the substituent
  - is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-
  - 4alkyloxy,
  - 30 (o) substituted thiazolyl C<sub>1</sub>-6alkyl wherein the substituent is
  - hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
  - (p) substituted oxazolyl C<sub>1</sub>-6alkyl wherein the substituent is
  - hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy;
- and



Y is  wherein R<sub>6</sub> and R<sub>7</sub> are each individually  
5 selected from the group consisting of:

- (a) H,
- (b) C<sub>1-10</sub>alkyl,
- (c) aryl, heteroaryl, arylC<sub>1-6</sub>alkyl or heteroarylC<sub>1-6</sub>alkyl,  
10 wherein the aryl or heteroaryl group is defined above.

### DETAILED DESCRIPTION OF THE INVENTION

The invention encompasses compounds of Formula I  
and contains several preferred subclasses as described below.

15 A preferred class is a compound according to Formula I  
wherein Ar is an aryl or heteroaryl group selected from the group  
consisting of:

- (1) phenyl,
- (2) naphthyl,
- 20 (3) pyridyl,
- (4) furyl,
- (5) thienyl,
- (6) isothiazolyl,
- (7) isoxazolyl,
- 25 (8) thiazolyl,
- (9) oxazolyl,


which can be mono- or di-substituted with the substituents  
independently selected from C<sub>1-6</sub>alkyl, halo, hydroxy, C<sub>1-6</sub>  
alkylcarbonyl, and aryl or heteroaryl are as defined above.  
30


A preferred embodiment of this class is a compound  
according to Formula I furthermore wherein R<sub>1</sub> is OH, OC<sub>1-6</sub>alkyl,  
OCH<sub>2</sub>phenyl.


Another class of preferred compounds is a compound according to Formula I wherein R<sub>2</sub> is hydrogen or mono- or di-substituted C<sub>1</sub>-6alkyl or C<sub>1</sub>-6 alkoxy, wherein the substituents are independently selected from the group consisting of:


- 5           (a) hydrogen,  
          (b) C<sub>1</sub>-6alkoxy,  
          (c) aryl or heteroaryl which is selected from the group consisting of
- 10           (1) phenyl,  
          (2) naphthyl,  
          (3) pyridyl,  
          (4) thienyl,  
          (5) imidazolyl,  
          (6) benzimidazolyl,  
15           (7) benzofuryl,  
          (8) benzothienyl,  
          (9) indolyl,  
          (10) isoxazolyl,  
          (11) thiazolyl,  
20           (12) oxazolyl,  
          (13) benzthiazolyl, and  
          (14) benzoxazolyl,


- 25           which can be mono- or di-substituted with substituents independently selected from C<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkyloxy, halo, hydroxy, amino, C<sub>1</sub>-6alkylamino, aminoC<sub>1</sub>-6alkyl, carboxyl, carboxylC<sub>1</sub>-6alkyl, and C<sub>1</sub>-6alkylcarbonyl;
- (d) aryloxy or heteroaryloxy wherein the aryl or heteroaryl groups are defined as in item (c) above;
- 30           (e) aroyl or heteroaroyl wherein the aryl or heteroaryl groups are defined as in item (c) above;

- (f)  wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; aryl or heteroaryl and mono and di-substituted aryl or heteroaryl as defined above in item (c); or substituted C<sub>1-6</sub>alkyl wherein the substituent is selected from hydroxy, halo, and aryl or heteroaryl, or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen and carbon atoms to which they are attached, to form a lactam or benzolactam ring wherein the lactam portion thereof is a ring of 5, 6, 7, or 8 atoms, said lactam or benzolactam containing a single hetero atom;


- (g)  wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; C<sub>1-6</sub> alkyl; aryl or heteroaryl as defined above in item (c);

- (h)  wherein R<sub>a</sub>, R<sub>b</sub>, and R<sub>c</sub> are each independently hydrogen; C<sub>1-6</sub> alkyl; aryl or heteroaryl as defined above in item (c);

- (i)  wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; C<sub>1-6</sub> alkyl; aryl or heteroaryl as defined above in item (c);



- (j) wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above in item (c); or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms and said nitrogen atom;



- (k) wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above in item (c); or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms and said nitrogen atom.


An another class is a compound according to Formula I wherein  $\text{R}_3$  is substituted C1-10alkylamino wherein the substituents are selected from the group consisting of:

- (1) hydrogen,
- (2) aminocarbonyl,
- (3) mono- or di-C1-6alkyl aminocarbonyl,
- (4) aryl or heteroaryl, wherein the aryl or heteroaryl group is selected from the group consisting of:
  - (a) phenyl,
  - (b) naphthyl,
  - (c) pyridyl,
  - (d) thienyl,
  - (e) imidazolyl,
  - (f) benzimidazolyl,
  - (g) benzofuryl,

- (h) benzothienyl,
- (i) indolyl,
- (j) isoxazolyl,
- (k) thiazolyl,
- (l) oxazolyl,
- (m) benzthiazolyl, and
- (n) benzoxazolyl,

which can be mono- or di-substituted with substituents independently selected from phenyl, C<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkyloxy, halo, hydroxy, amino, C<sub>1</sub>-6alkylamino, aminoC<sub>1</sub>-6alkyl, carboxyl, carboxylC<sub>1</sub>-6alkyl, or C<sub>1</sub>-6alkylcarbonyl.

A further preferred class is a compound according to Formula I wherein R<sub>3</sub> is an amino acid derivative of Formula II



wherein R<sub>f</sub> is hydrogen and R<sub>g</sub> is selected from:

- (a) C<sub>1</sub>-6alkyl,
- (b) amino C<sub>1</sub>-6alkyl
- (c) aminocarbonyl C<sub>1</sub>-6alkyl,
- (d) mono- or di-C<sub>1</sub>-6alkyl amino C<sub>1</sub>-6alkyl,
- (e) guanidino C<sub>1</sub>-6alkyl,
- (f) substituted phenyl C<sub>2</sub>-6alkyl, wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (g) substituted imidazolyl C<sub>3</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (h) substituted pyridyl C<sub>2</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (i) substituted pyridylamino C<sub>3</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4 alkyloxy,

- (j) substituted thiazolyl C<sub>2-6</sub>alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1-4</sub> alkyl, or C<sub>1-4</sub>alkyloxy,
- (k) substituted oxazolyl C<sub>2-6</sub>alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1-4</sub> alkyl, or C<sub>1-4</sub>alkyloxy.


5

A subclass of this class is a compound according to Formula I wherein Formula II is an amino acid selected from alanine, valine, leucine, isoleucine,  $\alpha$ -tert-butyl glycine, norleucine, serine, threonine, asparagine, glutamine, lysine, homohistidine, arginine, homophenylalanine, homotyrosine, methionine, ornithine, homoserine, and citrulline.

10

A still further subclass is a compound according to Formula I, wherein Formula II,

15



selected from the group consisting of:

20

- (a) C<sub>1-10</sub>alkyl,
- (b) aryl, heteroaryl, arylC<sub>1-6</sub>alkyl or heteroarylC<sub>1-6</sub>alkyl, wherein the aryl or heteroaryl group is selected from the group consisting of

25

30

- (1) phenyl,
- (2) naphthyl,
- (3) pyridyl,
- (4) thienyl,
- (5) imidazolyl,
- (6) benzimidazolyl,
- (7) benzofuryl,
- (8) benzothienyl,
- (9) indolyl,
- (10) isoxazolyl,
- (11) thiazolyl,
- (12) oxazolyl,


(13) benzthiazolyl, and

(14) benzoxazolyl,

and mono- and di-substituted aryl and heteroaryl groups as defined above in items (1) to (14) wherein the substituents are independently  
5 selected from phenyl, C<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkyloxy, halo, hydroxy, amino, C<sub>1</sub>-6alkylamino, aminoC<sub>1</sub>-6alkyl, carboxyl, carboxylC<sub>1</sub>-6alkyl, or C<sub>1</sub>-6alkylcarbonyl.

10 A preferred embodiment of this class is a compound according to Formula I wherein Ar is 4-propylphenyl or 4-biphenyl.

A still further subclass is a compound according to the Formula:



wherein:

p is 1-2;

Ar is 4-propylphenyl or 4-biphenyl;

30 R<sub>3</sub> is an amino derivative of Formula II wherein said amino acid is selected from alanine, valine, leucine, isoleucine, norleucine, α-tert-butyl glycine, serine, threonine, asparagine, glutamine, lysine, homohistidine, arginine, homophenylalanine, homotyrosine, methionine, ornithine, homoserine, or citrulline; wherein said Y in Formula II is NH-Phenyl.

Specific embodiments according to Formula I are the compounds:

- 5 (a) 1-(2-(4-Propylphenyl)ethyl)cyclopentane-1,3-dicarboxylic acid 1-(L-leucine N-phenylamide) amide;  
(b) 1-(2-(4-Propylphenyl)ethyl)cyclohexane-1,3-dicarboxylic acid 1-(L-leucine N-phenylamide) amide.

10 This invention also concerns a pharmaceutical composition and a method of treatment of stromelysin-mediated or implicated disorders or diseases (as described above) in a patient (which shall be defined to include man and/or mammalian animals raised in the dairy, meat, or fur industries or as pets) in need of such  
15 treatment comprising administration of the stromelysin inhibitors of Formula I as the active constituents.

Similarly, this invention also concerns pharmaceutical compositions and methods of treatment of collagenase mediated or implicated disorders or diseases (as described above) in a patient in  
20 need of such treatment comprising administration of the collagenase inhibitors of Formula I as the active constituents.


Similarly, this invention also concerns pharmaceutical compositions and methods of treatment of gelatinase-mediated or implicated disorders or diseases (as described above) in a patient in  
25 need of such treatment comprising administration of the gelatinase inhibitors of Formula I as the active constituents.

Moreover the invention also encompasses compositions, treatment, and method for co-administration of a compound of Formula I with a PMN elastase inhibitor such as those described in  
30 EP 0 337 549 which published on October 18, 1989, which is hereby incorporated by reference.


Compounds of the instant invention are conveniently prepared using the procedures described generally below in Scheme I and more explicitly described in the Example section thereafter.



SCHEME I



SCHEME I (CONT'D.)




The tert-butyl ester B of 4-(4-propylphenyl)butanoic acid A is prepared by reaction with tert-butyl alcohol in the presence of a condensing agent, eg. dicyclohexylcarbodiimide (DCC), and 4-dimethylaminopyridine (4-DMAP). This ester B is formylated with

tert-butoxybis(dimethylamino)methane to form t-butyl 2-formyl-4-(4-propylphenyl)butanoate C. According to the method described in the literature (P. L. Fuchs et al., *J. Org. Chem.* **1974**, *96*, 1607-9), the enolate of C is formed with sodium hydride and is reacted with  
 5 1-carboethoxycyclopropyl triphenylphosphonium tetrafluoroborate to form cyclopentene derivative D. Selective reduction of C is affected by hydrogenation in the presence of (tricyclohexylphosphine)(1,5-cyclooctadiene)(pyridine)iridium(I) hexafluorophosphate catalyst to afford the substituted cyclopentane  
 10 E. The ethyl ester in E is converted to the benzyl ester F by basic hydrolysis followed by esterification with benzyl alcohol. The t-butyl ester in F is deprotected by treatment with strong acid (eg. trifluoroacetic acid or hydrochloric acid) followed by condensation with an amino acid derivative, in this case, L-leucine N-phenylamide,  
 15 to form G. Catalytic hydrogenolysis of the benzyl ester affords the cyclopentane carboxylic acid H. Anyone skilled in the art would recognize that this scheme is representative of methodology that could be applied to the preparation of a variety of cyclopentane carboxylic acids of the type H by replacing A with other aryl  
 20 alkanoic acids and replacing L-leucine N-phenylamide with other amino acid derivatives.

25

30

SCHEME II



The t-butyl ester of cyclopentene D (prepared by methodology outline in Scheme I) is hydrolyzed in the presence of

strong acid (eg. trifluoroacetic acid or hydrochloric acid) followed  
by reaction with an amino acid derivative, in this case L-leucine N-  
phenylamide, to form I. Basic aqueous hydrolysis of I yields  
cyclopentene-carboxylic acid J. Catalytic hydrogenolysis of the  
5 double bond in J would yield the cyclopentane acid H. Anyone  
skilled in the art would recognize that this scheme is representative  
of methodology that could be applied to the preparation of a variety  
of cyclopentene carboxylic acids of the type J by replacing A from  
Scheme I with other aryl alkanoic acids and replacing L-leucine N-  
10 phenylamide with other amino acid derivatives.


15

20


25

30

SCHEME III



SCHEME III (CONT'D.)



The formyl ester C is reacted with methyl vinyl ketone in the presence of base followed by cyclization and elimination to


form cyclohexenone K. Catalytic hydrogenation affords cyclohexanone L which is acylated in the presence of strong base with methyl cyanoformate to form a ketoester. Reduction of the ketone, followed by sulfonylation of the resultant alcohol with mesyl chloride, and then base catalyzed elimination yields cyclohexene 1,3-diester M. Catalytic hydrogenation of the double bond in the presence of (tricyclohexyl-phosphine)(1,5-cyclooctadiene)-(pyridine)iridium(I) hexafluoro-phosphate catalyst affords cyclohexane 1,3-diester N. Methyl ester hydrolysis followed by condensation with benzyl alcohol yields diester O. t-Butyl ester hydrolysis followed by condensation with an amino acid derivative in the presence of benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP) affords cyclohexane P. Catalytic hydrogenolysis of the benzyl ester in P affords cyclohexane-carboxylic acid Q. Anyone skilled in the art would recognize that this scheme is representative of methodology that could be applied to the preparation of a variety of cyclohexane-carboxylic acids of the type Q by replacing A from Scheme I with other aryl alkanolic acids and replacing L-leucine N-phenylamide with other amino acid derivatives.

25

30



SCHEME IV



The t-butyl ester of cyclohexene M (prepared by methodology outlined in Scheme III) is hydrolyzed in the presence of


strong acid (eg. trifluoroacetic acid or hydrochloric acid) followed by reaction with an amino acid derivative, in this case L-leucine N-phenylamide, to form R. Basic aqueous hydrolysis of R yields cyclohexene-carboxylic acid S. Catalytic hydrogenolysis of the double bond in S yields the cyclohexane acid Q. Anyone skilled in the art would recognize that this scheme is representative of methodology that could be applied to the preparation of a variety of cyclohexene-carboxylic acids of the type S by replacing A from Scheme I with other aryl alkanoic acids and replacing L-leucine N-phenylamide with other amino acid derivatives.

The following examples are intended to illustrate the preparation of compounds of Formula I, and as such are not intended to limit the invention as set forth in the claims appended, thereto.

### EXAMPLE 1


1-(2-(4-Propylphenyl)ethyl)cyclopentane-1,3-dicarboxylic acid 1-(L-leucine phenylamide) amide

#### Step A:



**tert-Butyl 4-(4-propylphenyl)butanoate.** [The preparation of 4-(4-propylphenyl)butanoic acid, is described in the chemical literature: see L.I. Smith and C.-P. Lo, J. Am. Chem. Soc., 1948, 70, 2209-15 (Chem. Abstr. 42:7753c)].


4-(Dimethylamino)pyridine (265 mg, 2.17 mmol) was added to a solution of 4-(4-propylphenyl)butanoic acid (3.00 g, 14.5 mmol) and *tert*-butanol (4.10 mL, 3.22 g, 43.5 mmol) in dichloromethane (20 mL). The reaction was cooled to 0°C and a solution of 1,3-dicyclohexylcarbodiimide (3.59 g, 17.3 mmol) in dichloromethane (5.0 mL) was added over 10 min. The reaction was allowed to slowly warm to room temperature and was stirred for 24 h. Additional 1,3-dicyclohexylcarbodiimide (1.20 g, 5.82 mmol) was added and the reaction was stirred at room temperature of for 40 h. Water (3 mL) was added and the mixture was stirred for 15 min before being filtered. The precipitate was washed with hexane (25 mL) and the combined filtrate was washed with 2 N aq. HCl (25 mL), saturated aq. NaHCO<sub>3</sub> (25 mL), and saturated aq. NaCl (2 x 25 mL). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated. Purification by flash column chromatography on silica gel (110 g) eluting with 1 L of 5% Et<sub>2</sub>O/hexane gave 2.90 g (76% yield) of the title compound as a colorless oil. NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.10 (s, 4H), 2.60 (t, 2H, J = 7 Hz), 2.55 (t, 2H, J = 7 Hz), 2.23 (t, 2H, J = 7 Hz), 1.89 (quintet, 2H, J = 7 Hz), 1.62 (sextet, 2H, J = 7 Hz), 1.45 (s, 9H), 0.94 (t, 3H, J = 7 Hz).

Step B:

**tert-Butyl 2-formyl-4-(4-propylphenyl)butanoate.** A solution of 1.00 g (3.81 mmol) of *tert*-butyl 4-(4-propylphenyl)butanoate in *tert*-butoxybis(dimethylamino)methane (10.0 mL, 8.44 g, 48.4 mmol) was warmed in an oil bath at 120°C for 71 h. The solution was allowed to cool to room temperature before being poured into a 0°C solution containing water (12.5 mL) and concentrated aq. HCl (8.25 mL) dissolved in methanol (150 mL). After 5 min., the mixture was concentrated on a rotary evaporator and the residue was partitioned between EtOAc (75 mL) and 2 N aq HCl (40 mL). The organic layer was washed with saturated aq. NaCl (2 x 35 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), decanted, and evaporated. The residue was purified by flash column chromatography on silica gel (57 g), eluting with 500 mL of 5% Et<sub>2</sub>O/hexane followed by 250 mL of 10% Et<sub>2</sub>O/hexane. The starting ester (0.57 g, 57% recovery) was eluted before the title compound, isolated as a colorless oil (0.36 g, 32% yield).


NMR (400 MHz, CDCl<sub>3</sub>) indicated the presence of the product as a 2:1 mixture of the enol and aldehyde tautomers, respectively. Enol tautomer: δ 11.57 (d, 1H, J = 12 Hz), 7.09 (d, 2H, J = 8 Hz), 7.05 (d, 2H, J = 8 Hz), 6.90 (d, 1H, J = 12 Hz), 2.66 (t, 2H, J = 7 Hz), 2.55 (t, 2H, J = 7 Hz), 2.27 (t, 2H, J = 7 Hz), 1.62 (sextet, 2H, J = 7 Hz), 1.54 (s, 9H), 0.94 (t, 3H, J = 7 Hz). Aldehyde tautomer (partial data): d 9.68 (d, 1H, J = 2 Hz), 7.12-7.07 (m, 4H), 3.20 (td, 1H, J = 7, 2 Hz), 2.13 (q, 1H, J = 7 Hz), 1.50 (s, 9H).

Step C:



***tert*-Butyl 3-carbethoxy-1-(2-(4-propylphenyl)ethyl)-2-cyclopentene-1-carboxylate.** Sodium hydride (16 mg of 60% oil dispersion, 0.40 mmol) was added to a solution of 102 mg (0.35 mmol) of *tert*-butyl 2-formyl-4-(4-propylphenyl)butanoate in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (1.5 mL). After 20 min., 1-carboethoxycyclopropyl triphenylphosphonium tetrafluoroborate (193 mg, 0.42 mmol) was added to the pale yellow solution and stirring was continued overnight at room temperature. The solution was then partitioned between hexane (25 mL) and 2 N aq. HCl (25 mL). The organic layer was washed with water (15 mL) and saturated aq. NaCl (15 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), decanted, and evaporated. The residue was purified by flash column chromatography on silica gel (8 g), eluting with 100 mL of 5% EtOAc/hexane, to give the title compound as 52 mg (38% yield) of colorless oil.

NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.10 (d, 2H, *J* = 8 Hz), 7.04 (d, 3H, *J* = 8 Hz), 6.72 (t, 1H, *J* = 1.5 Hz), 4.21 (q, 2H, *J* = 7 Hz), 2.71-2.43 (m, 4H), 2.55 (t, 2H, *J* = 7 Hz), 2.06-1.88 (m, 4H), 1.62 (sextet, 2H, *J* = 7 Hz), 1.48 (s, 9 H), 1.31 (t, 3H, *J* = 7 Hz), 0.93 (t, 3H, *J* = 7 Hz).


Step D:

- 20 ***tert*-Butyl 3-carbethoxy-1-(2-(4-propylphenyl)ethyl)cyclopentane-1-carboxylate.** *tert*-Butyl 3-carbethoxy-1-(2-(4-propylphenyl)ethyl)-2-cyclopentene-1-carboxylate (69 mg, 0.179 mmol) was dissolved in dry dichloromethane (0.5 mL) in a flask connected to a hydrogen-filled balloon. (Tricyclohexylphosphine)(1,5-
- 25 cycloocta-diene)(pyridine)iridium(I) hexafluorophosphate (30 mg, 0.037 mmol) was dissolved in dry dichloromethane (1.0 mL), and 0.5 mL of the solution was added to the stirred hydrogenation reaction. After 2.5 h, the remaining catalyst solution was added and stirring was continued for an additional 2 h at room temperature.
- 30 The solution was then diluted with EtOAc (25 mL) and washed with 2 N aq. HCl (15 mL), saturated aq. NaHCO<sub>3</sub> (15 mL), and saturated aq. NaCl (15 mL). The solution was dried (Na<sub>2</sub>SO<sub>4</sub>), decanted, evaporated, and combined with crude product similarly obtained from 9 mg (0.02 mmol) of ester 4. Flash column chromatography twice on silica gel (10 g) eluting with 200 mL of 3% EtOAc/hexane

gave 47 mg (60% yield) of the more mobile stereoisomer of the title compound along with 24 mg (31% yield) of the less mobile stereoisomer.

NMR (400 MHz, CDCl<sub>3</sub>) of the more mobile product:  $\delta$  7.09 (d, 2H, J = 8 Hz), 7.06 (d, 2H, J = 8 Hz), 4.13 (q, 2H, J = 7 Hz), 2.89 (quintet, 1H, J = 7 Hz), 2.58-2.43 (m, 5 H), 2.16 (dt, 1 H, J = 12, 6 Hz), 2.05-1.83 (m, 4H), 1.71 (dd, 1H, J = 13, 9 Hz), 1.67-1.57 (m, 3H), 1.48 (s, 9H), 1.25 (t, 3H, J = 7 Hz), 0.93 (t, 3H, J = 7 Hz).

Step E:





**tert-Butyl 3-carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)-cyclopentane-1-carboxylate.** *tert*-Butyl 3-carbethoxy-1-(2-(4-propylphenyl)ethyl)cyclopentane-1-carboxylate (41 mg, 0.11 mmol) was dissolved in a mixture of THF (2.1 mL) and 95% ethanol (1.4 mL), and 0.5 N aq. LiOH (0.70 mL, 0.35 mmol) was added. After being stirred for 3 h at room temperature, the contents of the

reaction flask was partitioned between EtOAc (40 mL) and 2 N aq. HCl (20 mL). The organic layer was washed with saturated aqueous NaCl (20 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), decanted, and evaporated to give 39 mg of colorless film. This was combined with 5 mg of crude  
5 carboxylic acid obtained similarly from 6 mg of *tert*-butyl 3-carbethoxy-1-(2-(4-propylphenyl)-ethyl)cyclopentane-1-carboxylate and used directly in the esterification reaction.

Benzyl alcohol (0.025 mL, 26 mg, 0.24 mmol) and 4-(dimethylamino)pyridine (3 mg, 0.02 mmol) were added to a  
10 solution of the crude carboxylic acid (44 mg, 0.12 mmol) in dichloromethane (0.50 mL). A solution of 1,3-dicyclohexylcarbodiimide (32 mg, 0.16 mmol) in dichloromethane (0.10 mL) was added and the mixture was stirred 4 h at room temperature. Water (3 drops) and hexane (1.5 mL) were added and the mixture  
15 was filtered after 10 min. The precipitate was rinsed with hexane (5 mL) and the combined filtrate was diluted with hexane (15 mL) and washed with 2 N aq. HCl (10 mL) and saturated aq. NaCl (10 mL). The solution was dried (Na<sub>2</sub>SO<sub>4</sub>), decanted, and evaporated. The  
20 crude product was purified by flash column chromatography on silica gel (2.5 g) eluting with 100 mL of 5% EtOAc/hexane to give 52 mg (95% yield) of the title compound as a colorless oil. NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.39-7.29 (m, 5H), 7.09 (d, 2H, J = 8 Hz), 7.06 (d, 2H, J = 8 Hz), 5.12 (s, 2H), 2.96 (quintet, 1H, J = 8 Hz), 2.57-2.42 (m, 5 Hz), 2.16 (dt, 1H, J = 12, 6 Hz), 2.05-1.82 (m,  
25 4H), 1.74 (dd, 1H, J = 13, 9 Hz), 1.67-1.57 (m, 3H), 1.47 (s, 9H), 0.94 (t, 3H, J = 7 Hz).



Step F:

**3-Carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)cyclopentane-1-carboxylic acid (L-leucine phenylamide) amide.**

Trifluoroacetic acid (2.0 mL, 3.0 g, 26 mmol) was added to a stirred 0°C solution of 52 mg (0.12 mmol) of *tert*-butyl 3-carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)cyclopentane-1-carboxylate in dichloromethane (2.0 mL). After 2.5 h, toluene (2.0 mL) was added and the solution was evaporated to give crude carboxylic acid as 45 mg of pale yellow oil.

The crude carboxylic acid (45 mg, 0.11 mmol) was dried by evaporation of a toluene solution and then dissolved in DMF (0.40 mL). Molecular sieves (4 Å, 0.16 g) and N-methylmorpholine (0.027 mL, 25 mg, 0.25 mmol) were added. After 10 min., benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (60 mg, 0.14 mmol) was added and the solution was stirred for 40 min. before the addition of L-leucine anilide (28

mg, 0.14 mmol). After 1.5 h at room temperature, the solution was diluted with 20 mL of EtOAc and washed with 2 N aq. HCl (10 mL), saturated aq. NaHCO<sub>3</sub> (10 mL), and saturated aq. NaCl (10 mL).

The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), decanted, and evaporated.

5 Purification by flash column chromatography on silica gel (4.7 g) eluting with 100 mL of 3% EtOAc/dichloromethane gave 55 mg (83% yield) of the title compound as a colorless film.


NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers:

10  $\delta$  8.50 (s, 1 H), 7.47 (d, 2H, J = 7 Hz), 7.38-7.29 (m, 5H), 7.25 (t, 2H, J = 7 Hz), 7.06 (t, 1H, J = 7 Hz), 7.04 (d, 1H, J = 8 Hz), 7.02 (d, 1H, J = 8 Hz), 6.96 (d, 1H, J = 8 Hz), 6.93 (d, 1H, J = 8 Hz), 6.21 (d, 0.5H, J = 8 Hz), 6.16 (d, 0.5H, J = 8 Hz), 5.11 (s, 2H), 4.66 (bq, 1H, J = 7 Hz), 2.95 (quintet, 0.5 H, J = 8 Hz), 2.92 (quintet, 0.5H, J = 8 Hz), 2.60-2.37 (m, 5H), 2.16-2.06 (m, 1H), 2.00-1.54 (m, 11H), 0.99  
15 (d, 3H, J = 6 Hz), 0.97 (d, 1.5H, J = 6 Hz), 0.96 (d, 1.5 H, J = 6 Hz), 0.93 (t, 1.5H, J = 7 Hz), 0.92 (t, 1.5H, J = 7 Hz).

20

25

30

Step G:

**1-(2-(4-Propylphenyl)ethyl)cyclopentane-1,3-dicarboxylic acid 1-(L-leucine phenylamide) amide.** Catalyst (7 mg of 20% Pd(OH)<sub>2</sub>/C) was added to a solution of 3-carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)cyclopentane-1-carboxylic acid (leucine phenylamide) amide (55 mg, 0.94 mmol) in 95% ethanol (2.0 mL) and the mixture was stirred at room temperature for 2.5 h in a flask connected to a hydrogen-filled balloon. The mixture was centrifuged and the supernatant was filtered through a 0.45 micron membrane. The catalyst was rinsed with additional 95% ethanol and the combined filtrate was evaporated to give 46 mg (100% yield) of the title compound as a colorless brittle glass.


NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers:  $\delta$  8.82 (s, 0.5 H), 8.66 (s, 0.5H), 7.49 (d, 2H,  $J$  = 7 Hz), 7.25 (t, 2H,  $J$  = 7 Hz), 7.06 (bt, 1H,  $J$  = 7 Hz), 7.02 (d, 2H,  $J$  = 8 Hz), 6.95 (d,

1H, J = 8 Hz), 6.94 (d, 1H, J = 8 Hz), 6.76 (d, 0.5 H, J = 8 Hz), 6.70  
 (d, 0.5H, J = 8 Hz), 4.74 (bq, 1H, J = 7 Hz), 2.95-2.84 (m, 1H), 2.58  
 (dd, 0.5H, J = 13, 8 Hz), 2.54-2.36 (m, 2.5H), 2.50 (t, 2H, J = 7 Hz),  
 2.19-2.06 (m, 1H), 2.01-1.66 (m, 9H), 1.58 (sextet, 2H, J = 7 Hz),  
 0.98 (d, 3H, H = 6 Hz), 0.96 (d, 1.5 H, J = 6 Hz), 0.95 (1.5 H, J = 6  
 Hz), 0.91 (t, 3H, J = 7 Hz).

### EXAMPLE 2

1-(2-(4-Propylphenyl)ethyl)-2-cyclopentene-1,3-dicarboxylic acid 1-  
 (leucine phenylamide) amide

#### Step A:




**3-Carbethoxy-1-(2-(4-propylphenyl)ethyl)-2-cyclopentene-1-carboxylic acid (leucine phenylamide) amide.**

Trifluoroacetic acid (4.0 mL, 6.0 g, 52 mmol) was added to a stirred 0°C solution of 53 mg (0.14 mmol) of *tert*-butyl 3-carbethoxy-1-(2-(4-propylphenyl)-ethyl)-2-cyclopentene-1-carboxylate in dichloromethane (4.0 mL). After 2 h, toluene (5.0 mL) was added and the solution was evaporated to give crude carboxylic acid as 45 mg of colorless oil.

The crude carboxylic acid was dried by evaporation of a toluene solution and then dissolved in dimethylformamide (0.50 mL). Molecular sieves (4 Å, 0.23 g) and N-methylmorpholine (0.035 mL, 32 mg, 0.32 mmol) were added. After 10 min., benzotriazol-1-yloxytris-(dimethylamino)phosphonium hexafluorophosphate (80 mg, 0.18 mmol) was added and the solution was stirred for 40 min. before the addition of L-leucine anilide (37 mg, 0.18 mmol). After 2 h at room temperature, the solution was diluted with 30 mL of ethyl acetate and washed with 2 N aq. hydrochloric acid (20 mL), saturated aq. sodium bisulfate (20 mL), and saturated aq. sodium chloride (20 mL). The organic layer was dried over anhydrous sodium sulfate, decanted, and evaporated. Purification by flash column chromatography on silica gel (5 g) eluting with 150 mL of 1.5% ethyl acetate/dichloromethane followed by 50 mL of 3% ethyl acetate/dichloromethane gave 53 mg (73% yield) of the title compound as a colorless oil.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers: δ 8.51 (s, 0.5H), 8.47 (s, 0.5H), 7.50 (d, 1H, J = 7 Hz), 7.48 (d, 1H, J = 7 Hz), 7.27 (t, 2H, J = 7 Hz), 7.11-7.00 (m, 4H), 6.98 (d, 1H, J = 7 Hz), 6.78 (t, 0.5H, J = 1.5 Hz), 6.76 (t, 0.5H, J = 1.5 Hz), 6.06 (d, 0.6H, J = 7 Hz), 6.04 (d, 0.5H, J = 7 Hz), 4.68-4.60 (m, 1H), 4.23 (q, 2H, J = 6 Hz), 2.68-2.58 (m, 2H), 2.56-2.42 (m, 4H), 2.40-2.25 (m, 1H), 2.23-2.12 (m, 1H), 2.09-1.99 (m, 1H), 1.96-1.56 (m, 6H), 1.32 (t, 3H, J = 6 Hz), 0.99 (d, 1.5H, J = 6 Hz), 0.97 (d, 1.5H, J = 6 Hz), 0.96 (d, 1.5H, J = 6 Hz), 0.95 (d, 1.5H, J = 6 Hz), 0.93 (t, 1.5 H, J = 7 Hz), 0.92 (t, 1.5H, J = 7 Hz).

Step B:

25 **1-(2-(4-Propylphenyl)ethyl)-2-cyclopentene-1,3-dicarboxylic acid 1-(leucine phenylamide) amide. 3-Carboethoxy-1-(2-(4-propylphenyl)ethyl)-2-cyclopentene-1-carboxylic acid (leucine phenylamide) amide** (53 mg, 0.10 mmol) was dissolved in a mixture of tetrahydrofuran (1.9 mL) and 95% ethanol (1.3 mL), and 0.5 N aq. lithium hydroxide (0.65 mL, 0.33 mmol) was added. After being stirred for 24 h at room

30 temperature, the contents of the reaction flask was partitioned between ethyl acetate (30 mL) and 2 N aq. hydrochloric acid (15 mL). The organic layer was washed with saturated aqueous sodium chloride (15 mL), dried over anhydrous sodium sulfate, decanted, and evaporated to give 48 mg of crude product. Flash column

chromatography on silica gel (4 g) eluting with 20% ethyl acetate/hexane followed (50 mL) followed by 2% acetic acid/20% ethyl acetate/hexane (50 mL) yielded 45 mg (90% yield) of the title compound as a colorless brittle glass.


5 <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers: δ 9.13 (s, 0.5H), 8.85 (s, 0.5H), 7.76 (d, 0.5H, J = 7 Hz), 7.50 (d, 1H, J = 7.5 Hz), 7.43 (d, 1H, J = 7.5 Hz), 7.26 (t, 1H, J = 7.5 Hz), 7.21 (t, 1H, J = 7.5 Hz), 7.12-6.97 (m, 5.5H), 6.85 (s, 0.5H), 4.90 (q, 0.5H, J = 7 Hz), 4.81 (q, 0.5H, J = 7 Hz), 2.86-1.53  
10 (m, 15H), 1.00 (d, 1.5H, J = 6 Hz), 0.98-0.94 (m, 4.5H), 0.93 (t, 1.5H, J = 7 Hz), 0.91 (t, 1.5H, J = 7 Hz).

15

20

25

30

EXAMPLE 31-(2-(4-Propylphenyl)ethyl)cyclohexane-1,3-dicarboxylic acid 1-(leucine phenylamide) amideStep A:

**tert-Butyl 1-(2-(4-propylphenyl)ethyl)-2-cyclohexen-4-one-1-carboxylate.** Potassium *tert*-butoxide (12 mg, 0.11 mmol) was added to a solution of *tert*-butyl 2-formyl-4-(4-propylphenyl)butanoate (628 mg, 2.16 mmol) in 2.0 mL of *tert*-butanol. The solution was stirred at 25°C as methyl vinyl ketone (0.210 mL, 177 mg, 2.52 mmol) was added in portions over a 1 hr period. The mixture was stirred 1.5 h longer before being diluted with ethyl acetate (50 mL) and washed with 0.1N aq. hydrochloric acid (20 mL) followed by saturated aq. sodium chloride (20 mL). The



organic layer was dried anhydrous sodium sulfate, decanted, and evaporated to give 763 mg of crude intermediate product.

5 Molecular sieve pellets (4 Å, 0.64 g) were added to a solution of the crude product in toluene (4.0 mL). After addition of pyrrolidine (0.020 mL, 17 mg, 0.24 mmol) and acetic acid (0.040 mL, 42 mg, 0.70 mmol), the reaction was warmed in a 50°C oil bath for 45 min. The mixture was cooled to 25°C and diluted with ethyl acetate (40 mL). After washing with 2 *N* aq. hydrochloric acid (15 mL), saturated aq. sodium bicarbonate (15 mL), and saturated aq. sodium chloride (15 mL), the solution was dried over anhydrous sodium sulfate, decanted, and evaporated. Flash column chromatography on silica gel (25 g), eluting with 800 mL of 7% ethyl acetate/hexane gave 522 mg (71% yield) of the title compound as a colorless oil.


15 <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): δ 7.11 (d, 2H, *J* = 8 Hz), 7.07 (d, 2H, *J* = 8 Hz), 6.95 (d, 1H, *J* = 10 Hz), 6.00 (d, 1H, *J* = 10 Hz), 2.65-2.42 (m, 6H), 2.11-1.92 (m, 4H), 1.62 sextet, 2H, *J* = 7 Hz), 1.50 (s, 9H), 0.93 (t, 3H, *J* = 7 Hz).

20

25

30



Step B:



***tert*-Butyl 1-(2-(4-propylphenyl)ethyl)cyclohexan-4-one-1-carboxylate.** A solution of *tert*-butyl 1-(2-(4-propylphenyl)ethyl)-2-cyclohexen-4-one-1-carboxylate (519 mg, 1.52 mmol) in 95% ethanol (13 mL) was stirred with 10% Pd/C (50 mg) at room temperature in a round bottom flask connected to a hydrogen-filled balloon. After 6 h, the mixture was centrifuged. The supernatant was filtered through a 0.45 micron membrane and the catalyst was rinsed with additional 95% ethanol. Evaporation of the filtrate gave the title compound in quantitative yield.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): δ 7.11 (d, 2H, J = 8 Hz), 7.06 (d, 2H, J = 8 Hz), 2.58-2.42 (m, 8H), 2.38-2.30 (m, 2H), 1.88-1.82 (m, 2H), 1.70-1.57 (m, 4H), 1.53 (s, 9H), 0.93 (t, 3H, J = 7 Hz).

Step C:



**tert-Butyl 3-carbomethoxy-1-(2-(4-propylphenyl)ethyl)-3-cyclohexene-1-carboxylate.** A cyclohexane solution of lithium diisopropylamide mono(tetrahydrofuran) complex (0.54 mL, 1.5 M, 0.81 mmol) was added to tetrahydrofuran (4.0 mL) at -78°C. A solution of *tert*-butyl 1-(2-(4-propylphenyl)ethyl)cyclohexan-4-one-1-carboxylate (250 mg, 0.73 mmol) in tetrahydrofuran (0.40 mL) was added over 5 min, with additional tetrahydrofuran (0.20 mL) to rinse. After 45 min, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (0.10 mL) was added, followed 3 min later by methyl cyanoformate (0.070 mL, 75 mg, 0.88 mmol). The reaction mixture was stirred for 15 min at -78°C and then quenched by the addition of 1 N aq. hydrochloric acid (10 mL). The mixture was partitioned between ethyl acetate (80 mL) and additional 1 N aq. hydrochloric acid (30 mL). The ethyl acetate layer was washed with saturated aq. sodium chloride (40 mL), dried over anhydrous sodium sulfate,



decanted, and evaporated to give 0.36 g of yellow oil. Flash column chromatography on silica gel (10 g), eluting with 600 mL of 4% EtOAc gave 0.24 g of partially purified ketodiester intermediate.

5 The ketodiester (333 mg, 0.827 mmol) was dissolved in methanol (30 mL) and cooled to 0°C. Sodium borohydride (165 mg, 4.36 mmol) was added and the solution was stirred for 50 min before the addition of acetic acid (1.0 mL). After 10 min, most of the methanol was removed on a rotary evaporator and the residue was diluted with ethyl acetate (50 mL) and washed with 1 *N* aq.  
10 hydrochloric acid (50 mL), saturated aq. sodium bicarbonate (25 mL), and saturated aq. sodium chloride (25 mL). The organic solution was dried over anhydrous sodium sulfate, decanted, and evaporated to yield the crude hydroxydiester as 355 mg of colorless oil.

15 All of the crude hydroxydiester from the above step was dried by repeatedly dissolving in toluene (6 mL) and evaporating the solution. The residue was dissolved in dichloromethane (15 mL) and triethylamine (0.36 mL, 0.26 g, 2.6 mmol) was added. The solution was cooled to 0°C and methanesulfonyl chloride (0.095 mL, 140 mg,  
20 1.2 mmol) was added in portions over 45 min. After an additional 2 h at 0°C, the solution was diluted with ethyl acetate (35 mL) and washed with 2 *N* aq. hydrochloric acid (30 mL). The organic layer was partitioned with additional ethyl acetate (20 mL) and a mixture of saturated aq. sodium bicarbonate (30 mL) and saturated aq.  
25 sodium chloride (30 mL). The organic layer was washed with additional saturated aq. sodium chloride (2 x 20 mL), dried over anhydrous sodium sulfate, filtered, and evaporated to give the crude mesylate as 410 mg of colorless oil.

30 All of the crude mesylate from the above step was dried by repeatedly dissolving in toluene and evaporating the solution. The residue was dissolved in dichloromethane (55 mL) stirred at room temperature as 1,8-diazabicyclo[5.4.0]undec-7-ene (5.0 mL, 5.1 g, 33 mmol) was added. After 11 h, the solution was washed with 1 *N* aq. hydrochloric acid (50 mL), saturated aq. sodium


bicarbonate (25 mL), and saturated aq. sodium chloride (25 mL).  
 The aqueous layers were extracted successively with  
 dichloromethane (3 x 30 mL). The organic layers were combined,  
 dried over anhydrous sodium sulfate, decanted, and evaporated. The  
 residue was purified by flash column chromatography on silica gel  
 (40 g), eluting with 750 mL of 4% ethyl acetate/hexane to give the  
 title compound as 114 mg (29% overall yield) of colorless oil.  
<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): δ 7.10 (d, 2H, J = 8 Hz), 7.07 (d, 2H,  
 J = 8 Hz), 6.98-6.94 (m, 1H), 3.75 (s, 3H), 2.91 (d, 1H, J = 17 Hz),  
 2.62 (td, 1H, J = 12.5, 4.5 Hz), 2.55 (t, 2H, J = 7 Hz), 2.49 (td, 1H, J  
 = 12.5, 4.5 Hz), 2.30-2.23 (m, 2H), 2.12 (dd, 1H, J = 17, 2 Hz), 2.00  
 (dt, 1H, J = 13, 5 Hz), 1.91 (td, 1H, J = 12.5, 4.5 Hz), 1.76 (td, 1H, J  
 = 12.5, 4.5 Hz), 1.67-1.52 (m, 3H), 1.46 (s, 9H), 0.93 (t, 3H, J = 7  
 Hz).

Step D:

***tert*-Butyl 3-carbomethoxy-1-(2-(4-propylphenyl)ethyl)-cyclohexane-1-carboxylate.** A solution of (tricyclohexylphosphine)(1,5-cyclooctadiene)(pyridine)iridium(I) hexafluorophosphate (47 mg, 0.058 mmol) in dry dichloromethane (0.50 mL) was added to a round bottom flask containing *tert*-butyl 3-carbomethoxy-1-(2-(4-propylphenyl)ethyl)-3-cyclohexene-1-carboxylate (41 mg, 0.11 mmol) and connected to a hydrogen-filled balloon. After the stirring for 2 h at room temperature, additional catalyst (39 mg, 0.048 mmol) dissolved in dichloromethane (0.40 mL) was added. After another 3 h, the solution was diluted with ethyl acetate (15 mL) and washed with 2 N aq. hydrochloric acid (10 mL), saturated aq. sodium bicarbonate (10 mL), and saturated aq. sodium chloride (10 mL). The solution was dried over anhydrous sodium sulfate, decanted, evaporated, and combined with crude product similarly obtained from 8 mg (0.02 mmol) of the

unsaturated ester. Flash column chromatography three times on silica gel (5-8 g) eluting with 3-4% EtOAc/hexane gave 23 mg (47% yield) of the more mobile stereoisomer of the title compound. <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): δ 7.09 (d, 2H, J = 8 Hz), 7.04 (d, 2H, J = 8 Hz), 2.62-2.41 (m, 4 H), 2.54 (t, 2H, J = 7 Hz), 2.18 (bd, 1H, J = 12 Hz), 1.99-1.93 (m, 1H), 1.80 (td, 1H, J = 12.5, 4.5 Hz), 1.76-1.63 (m, 2 H), 1.62 (sextet, 2H, J = 7 Hz), 1.49 (s, 9H), 1.38-1.29 (m, 2H), 1.24 (t, 1H, J = 12 Hz), 1.10 (td, 1H, J = 12, 3.5 Hz), 0.93 (t, 3H, J = 7 Hz).

Step E:



***tert*-Butyl 3-carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)-cyclohexane-1-carboxylate.** *tert*-Butyl 3-carbomethoxy-1-(2-(4-propylphenyl)ethyl)cyclohexane-1-carboxylate (23 mg, 0.059 mmol) was dissolved in a mixture of THF (1.15 mL) and methanol (0.75

mL), and 0.5 N aq. lithium hydroxide (0.38 mL, 0.19 mmol) was added. After being stirred for 9 h at room temperature, the contents of the reaction flask was partitioned between ethyl acetate (20 mL) and 2 N aq. hydrochloric acid (10 mL). The organic layer was  
5 washed with saturated aqueous sodium chloride (10 mL), dried over anhydrous sodium sulfate, decanted, and evaporated to give 24 mg of crude product.


The crude carboxylic acid was dissolved in a dichloromethane solution (0.25 mL) containing benzyl alcohol (0.012 mL, 13  
10 mg, 0.12 mmol) and 4-(dimethylamino)pyridine (1.5 mg, 0.012 mmol). A solution of 1,3-dicyclohexylcarbodiimide (16 mg, 0.078 mmol) in dichloromethane (0.10 mL) was added. After 20 min., an additional portion of dichloromethane (0.10 mL) was added and the mixture was stirred 11 h at room temperature. Hexane (1.5 mL) and  
15 water (1 drop) were added. After 20 min., sodium sulfate was added and the mixture was filtered through a cotton plug with additional hexane (3 x 1.5 mL) to rinse. The filtrate was dried sodium sulfate, decanted, and evaporated. The crude product (36 mg) was purified by chromatography on a silica gel preparative thin layer  
20 chromatographic plate (20 x 20 x 0.2 cm) eluting with 4% ethyl acetate/hexane to give 22 mg (81% yield) of the title compound as a colorless oil.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): δ 7.39-7.30 (m, 5H), 7.09 (d, 2H, J = 8 Hz), 7.40 (d, 2H, J = 8 Hz), 5.14 (d, 1H, J = 12 Hz), 5.10 d, 1H, J = 12 Hz), 2.64-2.41 (m, 6H), 2.54 (t, 2H, J = 7 Hz), 2.18 (bd, 1H, J = 12 Hz), 2.02-1.96 (m, 1H), 1.80 (td, 1H, J = 12.5, 4.5 Hz), 1.76-1.65 (m, 2H), 1.62 sextet, 2H, J = 7 Hz), 1.48 (s, 9H), 1.39-1.30 (m, 2H), 1.27 (t, 1H, J = 12 Hz), 1.10 (td, 1H, J = 12, 3.5 Hz), 0.93 (t, 3H, J = 7 Hz).

30



Step F:




**3-Carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)cyclohexane-1-carboxylic acid (leucine phenylamide) amide.**

25 Trifluoroacetic acid (2.0 mL, 3.0 g, 26 mmol) was added to a stirred 0°C solution of 22 mg (0.047 mmol) of *tert*-butyl 3-carbobenzyloxy-1-(2-(4-propyl-phenyl)ethyl)cyclohexane-1-carboxylate in dichloromethane (2.0 mL). After 2.5 h, toluene (5.0 mL) was added and the solution was evaporated to give crude carboxylic acid as 20 mg of colorless film.

30 The crude carboxylic acid was dried by evaporation of a toluene solution and then dissolved in dimethylformamide (0.17 mL). Molecular sieves (4 Å, 68 mg) and N-methylmorpholine (0.012 mL, 11 mg, 0.11 mmol) were added. Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (26 mg, 0.059 mmol) was added and the solution was stirred for 40 min.

before the addition of L-leucine anilide (12 mg, 0.058 mmol). After 1.5 h at room temperature, the solution was diluted with 15 mL of ethyl acetate and washed with 2 N aq. hydrochloric acid (5 mL), saturated aq. sodium bicarbonate (5 mL), and saturated aq. sodium chloride (5 mL). The organic layer was dried over anhydrous sodium sulfate, decanted, and evaporated to give 35 mg of colorless film. Purification by flash column chromatography on silica gel (5 g) packed in dichloromethane, eluting with 100 mL of 1.5% ethyl acetate/dichloromethane gave 22 mg (79% yield) of the title compound as a colorless film.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers: δ 8.42 (s, 0.5H), 8.36 (s, 0.5H), 7.52-7.46 (m, 2H), 7.40-7.30 (m, 5H), 7.29-7.23 (m, 2H), 7.07 (t, 1H, J = 7 Hz), 7.02 (d, 1H, J = 8 Hz), 6.98 (d, 1H, J = 8 Hz), 6.92 (d, 1H, J = 8 Hz), 6.85 (d, 1H, J = 8 Hz), 6.15 (d, 0.5H, J = 7 Hz), 6.11 (d, 0.5H, J = 7 Hz), 5.14 (d, 0.5H, J = 12 Hz), 5.11 (s, 1H), 5.09 (d, 0.5H, J = 12 Hz), 4.70-4.61 (m, 1H), 2.62-2.32 (m, 6H), 2.14 (bd, 0.5H, J = 12 Hz), 2.08-1.99 (m, 1H), 1.99-1.92 (m, 0.5H), 1.84-1.52 (m, 8H), 1.43-1.22 (m, 4H), 0.99 (d, 1.5H, J = 6 Hz), 0.98-0.94 (m, 4.5H), 0.92 (t, 1.5H, J = 7 Hz), 0.91 (t, 1.5H, J = 7 Hz).

Step G:

25

30

**1-(2-(4-Propylphenyl)ethyl)cyclohexane-1,3-dicarboxylic acid 1-(leucine phenylamide) amide.** Catalyst (3.5 mg of 20%  $\text{Pd}(\text{OH})_2/\text{C}$ ) was added to a solution of 3-carbobenzyloxy-1-(2-(4-propylphenyl)ethyl)cyclohexane-1-carboxylic acid (leucine phenylamide) amide (22 mg, 0.037 mmol) in 95% ethanol (1.5 mL) and the mixture was stirred at room temperature for 2.5 h in a flask connected to a hydrogen-filled balloon. The mixture was centrifuged and the supernatant was filtered through a 0.45 micron membrane. The catalyst was rinsed with additional 95% ethanol and the combined filtrate was evaporated to give 18 mg (96% yield) of the title compound as a colorless glass.



$^1\text{H-NMR}$  (400 MHz,  $\text{CDCl}_3$ ) indicated a 1:1 mixture of diastereomers:  $\delta$  8.78 (s, 0.5H), 8.62 (s, 0.5H), 7.52 (d, 1H,  $J = 7$

Hz), 7.50 (d, 1H, J = 7 Hz), 7.43 (bs, 0.5H), 7.29 (t, 1H, J = 7 Hz),  
 7.27 (t, 1H, J = 7 Hz), 7.10 (t, 1H, J = 7 Hz), 7.04 (d, 1H, J = 8 Hz),  
 6.99 (t, 2H, J = 7 Hz), 6.90 (d, 1H, J = 8 Hz), 6.73 (d, 0.5 Hz), 4.83-  
 4.74 (m, 1H), 2.64-2.30 (m, 6H), 2.19 (d, 0.5H, J = 12 Hz), 2.09-  
 1.97 (m, 1H), 1.85-1.54 (m, 8.5H), 1.46-1.18 (m, 3.5H), 1.08 (bt,  
 0.5H, J = 12 Hz), 0.99 (d, 1.5 H, J = 6 Hz), 0.97 (d, 3H, J = 6 Hz),  
 0.96 (d, 1.5H, J = 6 Hz), 0.92 (t, 1H, J = 7 Hz), 0.91 (t, 1H, J = 7  
 Hz).

#### EXAMPLE 4

1-(2-(4-Propylphenyl)ethyl)-3-cyclohexene-1,3-dicarboxylic acid 1-  
 (leucine phenylamide) amide

##### Step A:




**3-Carbomethoxy-1-(2-(4-propylphenyl)ethyl)-3-cyclohexene-1-carboxylic acid (leucine phenylamide)**

amide. Trifluoroacetic acid (2.0 mL, 3.0 g, 26 mmol) was added to a stirred 0°C solution of 23 mg (0.060 mmol) of *tert*-butyl 3-carbomethoxy-1-(2-(4-propylphenyl)-ethyl)-3-cyclohexene-1-carboxylate in dichloromethane (2.5 mL). After 3 h, toluene (2.0 mL) was added and the solution was evaporated to give crude carboxylic acid as 21 mg of colorless film.

The crude carboxylic acid was dried by evaporation of a toluene solution and then dissolved in dimethylformamide (0.23 mL). Molecular sieves (4 Å, 0.10 g) and N-methylmorpholine (0.015 mL, 14 mg, 0.14 mmol) were added. After 10 min., benzotriazol-1-yloxytris-(dimethylamino)phosphonium hexafluorophosphate (34 mg, 0.077 mmol) was added and the solution was stirred for 40 min. before the addition of L-leucine anilide (16 mg, 0.078 mmol). After 2 h at room temperature, the solution was diluted with 20 mL of ethyl acetate and washed with 2 N aq. hydrochloric acid (10 mL), saturated aq. sodium bicarbonate (10 mL), and saturated aq. sodium chloride (10 mL). The organic layer was dried over anhydrous sodium sulfate, decanted, and evaporated. Chromatography on a silica gel preparative thin layer chromatographic plate (20 x 20 x 0.2 cm) eluting with 10% ethyl acetate/toluene gave 26 mg (84% yield) of the title compound as a colorless film.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers: δ 8.58 (s, 0.5H), 8.50 (s, 0.5H), 7.47 (d, 2H, J = 8 Hz), 7.27 (t, 1H, J = 8 Hz), 7.26 (t, 1H, J = 8 Hz), 7.11-6.87 (m, 6H), 6.16 (d, 0.5H, J = 7 Hz), 6.14 (d, 0.5H, J = 7 Hz), 4.68-4.60 (m, 1H), 3.75 (s, 1.5H), 3.74 (s, 1.5H), 2.89 (d, 0.5H, J = 16 Hz), 2.87 (d, 0.5H, J = 16 Hz), 2.63-2.22 (m, 7H), 2.08-1.90 (m, 2H), 1.84-1.54 (m, 7H), 0.99 (d, 1.5H, J = 6 Hz), 0.97 (d, 1.5H, J = 6 Hz), 0.96 (d, 1.5H, J = 6 Hz), 0.93 (t, 1.5H, J = 7 Hz), 0.92 (d, 1.5H, J = 6 Hz), 0.92 (t, 1.5H, J = 7 Hz).

Step B:

**1-(2-(4-Propylphenyl)ethyl)-3-cyclohexene-1,3-dicarboxylic acid 1-(leucine phenylamide) amide.** 3-Carbomethoxy-1-(2-(4-propylphenyl)ethyl)-3-cyclohexene-1-carboxylic acid (leucine phenylamide) amide (26 mg, 0.050 mmol) was dissolved in a mixture of THF (0.95 mL) and methanol (0.65 mL), and 0.5 N aq. lithium hydroxide (0.33 mL, 0.17 mmol) was added. After being stirred for 24 h at room temperature, the contents of the reaction flask was partitioned between ethyl acetate (30 mL) and 2 N aq. hydrochloric acid (15 mL). The organic layer was washed with saturated aqueous sodium chloride (15 mL), dried over sodium sulfate, decanted, and evaporated to give 26 mg of crude product. Flash column chromatography on silica gel (3 g) eluting with 20% ethyl acetate/hexane followed (50 mL) followed by

2% acetic acid/20% ethyl acetate/hexane (75 mL) yielded 13 mg (52% yield) of the title compound as a colorless film.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) indicated a 1:1 mixture of diastereomers:  $\delta$  8.95 (s, 0.5H), 8.89 (s, 0.5H), 7.70 (d, 0.5H, J = 7 Hz), 7.66 (d, 0.5H, J = 7 Hz), 7.52 (d, 1H, J = 7.5 Hz), 7.47 (d, 1H, J = 7.5 Hz), 7.25 (t, 2H, J = 7 Hz), 7.52-7.00 (m, 4H), 6.88 (d, 1H, J = 7 Hz), 6.72 (d, 1H, J = 7 Hz), 4.87-4.76 (m, 1H), 2.88-2.14 (m, 8H), 2.04-1.50 (m, 9H), 0.97 (d, 3H, J = 6 Hz), 0.95 (d, 3H, J = 6 Hz), 0.93 (t, 1.5H, J = 7 Hz), 0.90 (t, 1.5H, J = 7 Hz).

### EXAMPLE 5

#### Enzyme Inhibition Assays

##### **15 Inhibition of Human Fibroblast Stromelysin:**

Activation: Human recombinant stromelysin was purchased from Celltech (Slough, U.K.) as a proenzyme of 55kD in a buffer consisting of 20 mM Tris, 10 mM CaCl<sub>2</sub>, 0.05% Brij-35, and 0.2% NaN<sub>3</sub>, pH=7.5. The material was activated as described.(Harrison, R.K., Teahan, J., and Stein, R.L. A Semi-Continuous HPLC-Based Assay for Stromelysin. *Anal. Biochem.* **1989**, 180, 110-113). Briefly, to 1.0 mL of a 2.2  $\mu$ M solution of prostromelysin was added 20  $\mu$ L of a 1.0  $\mu$ M solution of trypsin in assay buffer (20 mM HEPES, 10mM CaCl<sub>2</sub>, 0.05% Brij-35, pH=7.5, [trypsin]<sub>final</sub> = 20 nM. The solution was incubated at 37°C for 30 minutes. The reaction was quenched by addition of a 50 fold molar excess of soybean trypsin inhibitor bound to agarose (Sigma), and the solution centrifuged to remove the trypsin:inhibitor complex.

**30 K<sub>i</sub> Determinations:** Stock solutions of inhibitors were prepared by dissolving the compounds in DMSO. The inhibitors were further diluted in assay buffer to eight different concentrations encompassing the approximate K<sub>i</sub>, and covering a 200 fold range. To 50  $\mu$ L of each of the inhibitor solutions was added 25  $\mu$ L of an 8 nM solution of trypsin-activated stromelysin, [DMSO] = 1.8%. The

solution was allowed to incubate for 4 h to reach equilibrium. To this solution was added 60  $\mu\text{L}$  of a 12.8  $\mu\text{M}$  solution of the substrate Arg-Pro-Lys-Pro-Leu-Ala-Phe-Trp-NH<sub>2</sub> ( $k_c/K_m = 12,000 \text{ M}^{-1}\text{s}^{-1}$ ) (Niedzwiecki, L., Teahan, J., Harrison, R.K., Stein, R.L. Substrate  
5 Specificity of the Human Metalloproteinase Stromelysin and the Development of Continuous, Fluorometric Assays. *Biochem.* **1992**, *31*, 12618-12623), and the reaction was allowed to proceed for 18 h. In the reaction  $[S] = 5.7 \mu\text{M}$  and  $[E] = 1.5 \text{ nM}$ . The reaction was  
10 quenched by addition of 50  $\mu\text{L}$  of 0.15M phosphoric acid, and 100  $\mu\text{L}$  of the reaction mixture was injected onto the HPLC. Since the reactions were run under first order conditions, ( $[S] \ll K_m$ ,  $K_m = 0.5 \text{ mM}$ ), the pseudo first order rate constant,  $k_{\text{obs}}$ , was determined for each of the inhibitor concentrations from the peak area  
15 corresponding to unreacted substrate in the inhibited sample, and the peak area for substrate at time = 0:

$$\ln \left( \frac{\text{area}_{\text{inhib}}}{\text{area}_{t=0}} \right) = -k_{\text{obs}}t$$

20

Values of  $K_i$  were determined from the ratio of the rate constants for inhibited and control sample (no inhibitor) plotted as a function of the inhibitor concentration, and fit to the following  
25 equation:

$$\frac{k_{\text{inhib}}}{k_{\text{control}}} = \frac{1}{1 + ([I] / K_i)}$$

30

### **Inhibition of Human Fibroblast Collagenase:**

Activation: Human fibroblast collagenase was purchased from Celltech (Slough, U.K.). The material was received as a proenzyme of 54 kD at a concentration of 1.2  $\mu\text{M}$  in a buffer consisting of 20 mM Tris, 5 mM CaCl<sub>2</sub>, 0.15 M NaCl, and 0.01% NaN<sub>3</sub>. The



material was activated with trypsin using the same procedure as for stromelysin, with the addition that the activation buffer contained 40 nM prostromelysin.

5 Ki Determinations. Stock solutions of inhibitors were prepared by dissolving the material in DMSO. The inhibitors were further diluted in assay buffer to eight different concentrations encompassing the approximate  $K_i$  and covering a 200 fold range. Final DMSO  
10 concentration was 2.8%. To 50  $\mu$ L of the inhibitor solutions was added 25  $\mu$ L of a 108 nM solution of trypsin activated collagenase. The solution was allowed to incubate for 4 h to reach equilibrium. To this solution was added 60  $\mu$ L of a 56  $\mu$ M solution of the substrate DNP-Pro-Leu-Gly-Leu-Trp-Ala-dArg-NH<sub>2</sub> ( $k_{cat}/K_m$  =  
15 270,000  $M^{-1}s^{-1}$ ) (Stack, M. S.; Gray, R. D. Comparison of Vertebrate Collagenase and Gelatinase Using a New Fluorogenic Substrate Peptide. *J. Biol. Chem.* 1989, 264, 4277-4281), and the reaction was allowed to proceed for 18 h. In the incubation mixture [S] = 25  $\mu$ M, [E] = 20 nM. The reaction was quenched by addition of  
20 50  $\mu$ L of 0.15 M phosphoric acid. The reaction mixture was injected onto the HPLC. The calculation of  $K_i$  was the same as for stromelysin.

#### **Inhibition of Human Gelatinase A:**

25 Activation: Human 72kD gelatinase was purchased from Celltech (Slough, U.K.) as a proenzyme at a concentration of 1.5  $\mu$ M in a buffer consisting of 20 mM Tris, 5mM CaCl<sub>2</sub> 150 mM NaCl, 0.01% brij, 0.02% NaN<sub>3</sub>, pH = 7.5. The proenzyme was activated by incubation of 500  $\mu$ L proenzyme with 50  $\mu$ L of a 11 mM solution  
30 aminophenyl mercuric acetate in NaOH (pH=11) at 25°C for 120 min.

Ki Determinations. Determination of  $K_i$  values for gelatinase A were identical to that of collagenase and stromelysin with the

exception that incubation of the enzyme-inhibitor mixture with the substrate was performed for only 2 h.  $[S] = 25 \mu M$ ,  $[E] = 20 nM$ .

5           This invention also relates to a method of treatment for patients (including man and/or mammalian animals raised in the dairy, meat, or fur industries or as pets) suffering from disorders or diseases which can be attributed to stromelysin as previously described, and more specifically, a method of treatment involving the administration of the matrix metalloendoproteinase inhibitors of  
10       Formula I as the active constituents.

          Accordingly, the compounds of Formula I can be used among other things in the treatment of osteoarthritis and rheumatoid arthritis, and in diseases and indications resulting from the over-expression of these matrix metalloendoproteinases such as found in  
15       certain metastatic tumor cell lines.

          For the treatment of rheumatoid arthritis, osteoarthritis, and in diseases and indications resulting from the over-expression of matrix metalloendoproteinases such as found in certain metastatic tumor cell lines or other diseases mediated by the matrix  
20       metalloendo-proteinases, the compounds of Formula I may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections,  
25       intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats, etc., the compounds of the invention are effective in the treatment of humans.

          The pharmaceutical compositions containing the active  
30       ing. lient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of

pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable  
5 preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate;  
10 granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the  
15 gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in the U.S. Patents 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic  
20 tablets for control release.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is  
25 mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example  
30 sodium carboxymethylcellulose, methylcellulose, hydroxypropyl-methyl-cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for

example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such  
5 as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more  
10 preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin.  
15 The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

20 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are  
25 exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be  
30 a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from

fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

5

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

10

15

20

25

The compounds of Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

30

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of Formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

Dosage levels of the order of from about 0.05 mg to about 140 mg per kilogram of body weight per day are useful in the

5 treatment of the above- indicated conditions (about 2.5 mg to about 7  
gms. per patient per day). For example, inflammation may be  
effectively treated by the administration of from about 0.01 to 50 mg  
of the compound per kilogram of body weight per day (about 0.5 mg  
to about 3.5 gms per patient per day).

10 The amount of active ingredient that may be combined  
with the carrier materials to produce a single dosage form will vary  
depending upon the host treated and the particular mode of  
administration. For example, a formulation intended for the oral  
administration of humans may contain from 0.5 mg to 5 gm of active  
agent compounded with an appropriate and convenient amount of  
carrier material which may vary from about 5 to about 95 percent of  
the total composition. Dosage unit forms will generally contain  
between from about 1 mg to about 500 mg of an active ingredient.

15 It will be understood, however, that the specific dose  
level for any particular patient will depend upon a variety of factors  
including the activity of the specific compound employed, the age,  
body weight, general health, sex, diet, time of administration, route  
of administration, rate of excretion, drug combination and the  
severity of the particular disease undergoing therapy.


20 The following Examples are intended to illustrate the  
preparation of compounds of Formula I, and as such are not  
intended to limit the invention as set forth in the claims appended,  
thereto.

25

30

WHAT IS CLAIMED IS:

1. A Compound of Formula I:



15 wherein:

n is 1 - 4;

p is 1-2;

20

Ar is an aryl or heteroaryl group selected from the group consisting of:

- (1) phenyl,
- (2) naphthyl,
- 25 (3) pyridyl,
- (4) pyrrol,
- (5) furyl,
- (6) thienyl,
- (7) isothiazolyl,
- 30 (8) imidazolyl,
- (9) benzimidazolyl,
- (10) tetrazolyl,
- (11) pyrazinyl,
- (12) pyrimidyl,
- (13) quinolyl,

- 5 (14) isoquinolyl,  
(15) benzofuryl,  
(16) isobenzofuryl,  
(17) benzothienyl,  
(18) pyrazolyl,  
(19) indolyl,  
(20) isoindolyl,  
(21) purinyl,  
10 (22) carboxazolyl,  
(23) isoxazolyl,  
(24) thiazolyl,  
(25) oxazolyl,  
(26) benzthiazolyl, and  
(27) benzoxazolyl,

15 which can be mono- or di-substituted with substituents  
independently selected from C<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkyloxy, halo,  
hydroxy, amino, C<sub>1</sub>-6alkylamino, aminoC<sub>1</sub>-6alkyl, carboxyl,  
carboxylC<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkylcarbonyl, aryl or heteroaryl as defined  
above;

20

R<sub>1</sub> is NHOH, OH, OC<sub>1</sub>-10alkyl, OCH(C<sub>1</sub>-10alkyl)OC(O)C<sub>1</sub>-10alkyl,  
OCH<sub>2</sub>phenyl, wherein the phenyl group may be substituted  
with hydrogen, carboxy, carboxyC<sub>1</sub>-3alkyl, aminocarbonyl,  
C<sub>1</sub>-6 alkylaminocarbonyl;

25


R<sub>2</sub> is hydrogen or a mono- or di-substituted C<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkoxy,  
or C<sub>2</sub>-6alkenyl wherein the substituents are independently  
selected from the group consisting of:

- 30 (a) hydrogen,  
(b) carboxy,  
(c) aminocarbonyl,  
(d) C<sub>1</sub>-6alkoxy,  
(e) C<sub>1</sub>-6alkylcarbonyl,  
(f) aryl or heteroaryl, as defined above;



- (g) aryloxy or heteroaryloxy wherein the aryl or heteroaryl groups are defined above;
- (h) aroyl or heteroaroyl wherein the aryl or heteroaryl groups are defined above;

5




- (i) wherein  $R_a$  and  $R_b$  are each independently hydrogen; aryl or heteroaryl and mono and di-substituted aryl or heteroaryl as defined above; or substituted  $C_{1-6}$ alkyl wherein the substituent is selected from hydroxy, halo, and aryl or heteroaryl, or wherein  $R_a$  and  $R_b$  are joined together with the nitrogen and carbon atoms to which they are attached, to form a lactam or benzolactam ring wherein the lactam portion thereof is a ring of 5, 6, 7, or 8 atoms, said lactam or benzolactam containing a single hetero atom;

10

15

20



- (j) wherein  $R_a$  and  $R_b$  are each independently hydrogen; aryl or heteroaryl and mono and di-substituted aryl or heteroaryl as defined above ; or substituted  $C_{1-6}$ alkyl wherein the substituent is selected from hydroxy, halo, and aryl, or wherein  $R_a$  and  $R_b$  are joined together with the nitrogen and carbon atoms to which they are attached to form a cyclic imide, or is benzofused onto a cyclic imide, wherein the cyclic imide portion thereof is a ring of 5, 6, 7, or 8 atoms, said cyclic imide containing a single hetero atom;

25

30

- (k) amino or mono- or di-substituted amino wherein the substituents are independently selected from C1-6 alkyl and aryl or heteroaryl as defined above;

5

- (l) 
$$\begin{array}{c} \text{O} \\ \parallel \\ \text{---N---C---O---R}_b \\ | \\ \text{R}_a \end{array}$$
 wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above ; or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen and oxygen atoms to which they are attached, to form a cyclic-carbamate, or a benzofused cyclic-carbamate, wherein the carbamate ring contains 5, 6, 7, or 8 atoms, said ring containing the two heteroatoms;

10

15

- (m) 
$$\begin{array}{c} \text{O} \\ \parallel \\ \text{---N---C---N---R}_b \\ | \quad | \\ \text{R}_a \quad \text{R}_c \end{array}$$
 wherein R<sub>a</sub>, R<sub>b</sub>, and R<sub>c</sub> are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen atoms to which they are attached to form a cyclic-urea or benzofused cyclic-urea, said urea ring containing 5, 6, 7, or 8 atoms and said two nitrogens;

20

25

- (n) 
$$\begin{array}{c} \text{O} \\ \parallel \\ \text{---N---S---R}_b \\ | \quad \parallel \\ \text{R}_a \quad \text{O} \end{array}$$
 wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein R<sub>a</sub> and R<sub>b</sub> are joined together with the nitrogen and sulfur atoms to which they are attached, to form a cyclic-sulfonamide or benzofused cyclic-sulfonamide ring, said sulfonamide ring containing 5, 6, 7, or 8 atoms and said heteroatoms;

30

- (o) 
$$\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{N}-\text{R}_b \\ | \\ \text{R}_a \end{array}$$
 wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms and said nitrogen atom;


- (p) 
$$\begin{array}{c} \text{O} \\ \parallel \\ -\text{S}-\text{N}-\text{R}_b \\ \parallel \quad | \\ \text{O} \quad \text{R}_a \end{array}$$
 wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C1-6 alkyl; aryl or heteroaryl as defined above; or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms and said nitrogen atom;

$\text{R}_3$  is

- (a) mono- or di-substituted C1-10alkylamino, or mono- or di-substituted di-C1-10alkyl amino wherein the substituents are selected from the group consisting of:
- (1) hydrogen,
  - (2) mercapto,
  - (3) hydroxy,
  - (4) carboxy,
  - (5) amino,
  - (6) aminocarbonyl,
  - (7) mono- or di-C1-6alkyl amino,
  - (8) mono- or di-C1-6alkyl aminocarbonyl,
  - (9) guanidino,
  - (10) aryl or heteroaryl, as defined above;

(b) an amino acid derivative of Formula II


5



wherein R<sub>f</sub> and R<sub>g</sub> are individually selected from:

10

- (a) hydrogen,
- (b) C<sub>1</sub>-6alkyl,
- (c) mercapto C<sub>1</sub>-6alkyl,
- (d) hydroxy C<sub>1</sub>-6alkyl,
- (e) carboxy C<sub>1</sub>-6alkyl,
- 15 (f) amino C<sub>1</sub>-6alkyl
- (g) aminocarbonyl C<sub>1</sub>-6alkyl,
- (h) mono- or di-C<sub>1</sub>-6alkyl amino C<sub>1</sub>-6alkyl,
- (i) guanidino C<sub>1</sub>-6alkyl,
- (j) substituted phenyl C<sub>1</sub>-6alkyl, wherein the substituent is
- 20 (k) substituted indolyl C<sub>1</sub>-6alkyl, wherein the substituent is
- (l) substituted imidazolyl C<sub>2</sub>-6alkyl wherein the substituent is
- (m) substituted pyridyl C<sub>1</sub>-6alkyl wherein the substituent is
- 25 (n) substituted pyridylamino C<sub>1</sub>-6alkyl wherein the substituent
- is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (o) substituted thiazolyl C<sub>1</sub>-6alkyl wherein the substituent is
- 30 (p) substituted oxazolyl C<sub>1</sub>-6alkyl wherein the substituent is
- hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy;
- and

Y is  wherein R<sub>6</sub> and R<sub>7</sub> are each individually  
5 selected from the group consisting of:

- (a) H,
- (b) C<sub>1-10</sub>alkyl,
- (c) aryl, heteroaryl, arylC<sub>1-6</sub>alkyl or heteroarylC<sub>1-6</sub>alkyl,  
10 wherein the aryl or heteroaryl group is defined above.

2. A compound according to Claim 1 wherein  
Ar is an aryl or heteroaryl group selected from the group consisting  
of:

- (1) phenyl,
- (2) naphthyl,
- (3) pyridyl,
- (4) furyl,
- (5) thienyl,
- (6) isothiazolyl,
- (7) isoxazolyl,
- (8) thiazolyl,
- (9) oxazolyl,

which can be mono- or di-substituted with the substituents  
independently selected from C<sub>1-6</sub>alkyl, halo, hydroxy, C<sub>1-6</sub>  
25 alkylcarbonyl, and aryl or heteroaryl as defined above in Claim 1;

3. A compound according to Claim 1 wherein R<sub>1</sub> is  
OH, OC<sub>1-6</sub>alkyl, OCH<sub>2</sub>phenyl.

4. A compound according to Claim 1 wherein R<sub>2</sub> is  
hydrogen or mono- or di-substituted C<sub>1-6</sub>alkyl or C<sub>1-6</sub> alkoxy,  
wherein the substituents are independently selected from the group  
consisting of:


- (a) hydrogen,

- (b) C<sub>1-6</sub>alkoxy,  
 (c) aryl or heteroaryl which is selected from the group consisting of

- (1) phenyl,
- (2) naphthyl,
- (3) pyridyl,
- (4) thienyl,
- (5) imidazolyl,
- (6) benzimidazolyl,
- (7) benzofuryl,
- (8) benzothienyl,
- (9) indolyl,
- (10) isoxazolyl,
- (11) thiazolyl,
- (12) oxazolyl,
- (13) benzthiazolyl, and
- (14) benzoxazolyl,

which can be mono- or di-substituted with substituents independently selected from C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, halo, hydroxy, amino, C<sub>1-6</sub>alkylamino, aminoC<sub>1-6</sub>alkyl, carboxyl, carboxylC<sub>1-6</sub>alkyl, and C<sub>1-6</sub>alkylcarbonyl;

- (d) aryloxy or heteroaryloxy wherein the aryl or heteroaryl groups are defined as in item (c) above;  
 (e) aroyl or heteroaroyl wherein the aryl or heteroaryl groups are defined as in item (c) above;




- (f) wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen; aryl or heteroaryl and mono and di-substituted aryl or heteroaryl as defined above in item (c); or substituted C<sub>1-6</sub>alkyl wherein the substituent is selected from hydroxy, halo, and aryl or heteroaryl, or wherein R<sub>a</sub>


and R<sub>b</sub> are joined together with the nitrogen and carbon atoms to which they are attached, to form a lactam or benzolactam ring wherein the lactam portion thereof is a ring of 5, 6, 7, or 8 atoms, said lactam or benzolactam containing a single hetero atom;

5


10



15





20



25

30





- (k) wherein  $\text{R}_a$  and  $\text{R}_b$  are each independently hydrogen; C<sub>1</sub>-6 alkyl; aryl or heteroaryl as defined above in item (c); or wherein  $\text{R}_a$  and  $\text{R}_b$  are joined together with the nitrogen atom to which they are attached, to form a heterocycle or benzofused heterocycle ring, said ring containing 5, 6, 7, or 8 atoms, and said nitrogen atom.

5. A compound according to Claim 1 wherein  $\text{R}_3$  is substituted C<sub>1</sub>-10alkylamino wherein the substituents are selected from the group consisting of:


- (1) hydrogen,
- (2) aminocarbonyl,
- (3) mono- or di-C<sub>1</sub>-6alkyl aminocarbonyl,
- (4) aryl or heteroaryl, wherein the aryl or heteroaryl group is selected from the group consisting of:
  - (a) phenyl,
  - (b) naphthyl,
  - (c) pyridyl,
  - (d) thienyl,
  - (e) imidazolyl,
  - (f) benzimidazolyl,
  - (g) benzofuryl,
  - (h) benzothienyl,
  - (i) indolyl,
  - (j) isoxazolyl,
  - (k) thiazolyl,
  - (l) oxazolyl,
  - (m) benzthiazolyl, and
  - (n) benzoxazolyl,

which can be mono- or di-substituted with substituents independently selected from phenyl, C<sub>1</sub>-6alkyl, C<sub>1</sub>-6alkyloxy, halo,



hydroxy, amino, C<sub>1</sub>-6alkylamino, aminoC<sub>1</sub>-6alkyl, carboxyl, carboxylC<sub>1</sub>-6alkyl, or C<sub>1</sub>-6alkylcarbonyl.

6. A compound according to Claim 1 wherein R<sub>3</sub> is an amino acid derivative of Formula II




wherein R<sub>f</sub> is hydrogen and R<sub>g</sub> is selected from:

- (a) C<sub>1</sub>-6alkyl,
- (b) amino C<sub>1</sub>-6alkyl
- (c) aminocarbonyl C<sub>1</sub>-6alkyl,
- (d) mono- or di-C<sub>1</sub>-6alkyl amino C<sub>1</sub>-6alkyl,
- (e) guanidino C<sub>1</sub>-6alkyl,
- (f) substituted phenyl C<sub>2</sub>-6alkyl, wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (g) substituted imidazolyl C<sub>3</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (h) substituted pyridyl C<sub>2</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (i) substituted pyridylamino C<sub>3</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (j) substituted thiazolyl C<sub>2</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy,
- (k) substituted oxazolyl C<sub>2</sub>-6alkyl wherein the substituent is hydrogen, hydroxy, carboxy, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4alkyloxy.

7. A compound according to Claim 6 wherein Formula II is an amino acid selected from alanine, valine, leucine, isoleucine, α-tert-butyl glycine, norleucine, serine, threonine, asparagine, glutamine, lysine, homohistidine, arginine,

homophenylalanine, homotyrosine, methionine, ornithine, homoserine, and citrulline.

8. A compound according to Claim 6 wherein




selected from the group consisting of:

- (a)  $C_{1-10}$ alkyl,
- (b) aryl, heteroaryl, aryl $C_{1-6}$ alkyl or heteroaryl $C_{1-6}$ alkyl, wherein the aryl or heteroaryl group is selected from the group consisting of
  - (1) phenyl,
  - (2) naphthyl,
  - (3) pyridyl,
  - (4) thienyl,
  - (5) imidazolyl,
  - (6) benzimidazolyl,
  - (7) benzofuryl,
  - (8) benzothieryl,
  - (9) indolyl,
  - (10) isoxazolyl,
  - (11) thiazolyl,
  - (12) oxazolyl,
  - (13) benzthiazolyl, and
  - (14) benzoxazolyl,

and mono- and di-substituted aryl and heteroaryl groups as defined above in items (1) to (14) wherein the substituents are independently selected from phenyl,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, halo, hydroxy, amino,  $C_{1-6}$ alkylamino, amino $C_{1-6}$ alkyl, carboxyl, carboxyl $C_{1-6}$ alkyl, or  $C_{1-6}$ alkylcarbonyl.

9. A compound according to Claim 1 wherein Ar is 4-propylphenyl or 4-biphenyl.

10. A compound according to Claim 1 of the  
Formula:



wherein:

p is 1-2;

Ar is 4-propylphenyl or 4-biphenyl;

R<sub>3</sub> is an amino derivative of Formula II wherein said amino acid is selected from alanine, valine, leucine, isoleucine, norleucine, α-tert-butyl glycine, serine, threonine, asparagine, glutamine, lysine, homohistidine, arginine, homophenylalanine, homotyrosine, methionine, ornithine, homoserine, or citrulline; wherein said Y in Formula II is NH-Phenyl.

11. A compound according to Claim 10 which is:

- (a) 1-(2-(4-Propylphenyl)ethyl)cyclopentane-1,3-dicarboxylic acid 1-(L-leucine N-phenylamide) amide, or
- (b) 1-(2-(4-Propylphenyl)ethyl)cyclohexane-1,3-dicarboxylic acid 1-(L-leucine N-phenylamide) amide.

12. A pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a compound according to Claim 1.

5

13. A pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a compound according to Claim 11.

10

14. A method of treating matrix metalloendoproteinase-mediated disease comprising administering to a subject in need thereof, a therapeutically effective amount of a compound according to Claim 1.

15

15. A method of treating matrix metalloendoproteinase-mediated disease comprising administering to a subject in need thereof, a therapeutically effective amount of a compound according to Claim 11.

20

16. A method of treating degenerative disease comprising the administration to a subject in need thereof a therapeutically effective amount of a compound according to Claim 1.

25

17. A method of treating degenerative disease comprising administering to a subject in need thereof a therapeutically effective amount of a compound according to Claim 11.

18. The use of a compound as claimed in any of claims 1 to 11 for the manufacture of a medicament.

**Patents Act 1977**  
**Examiner's report to the Comptroller under Section 17**  
**(The Search report)**

Application number  
GB 9420114.2

- 84 -

**Relevant Technical Fields**

(i) UK Cl (Ed.N) C2C (CKK, CKN, CKS, CK2)

(ii) Int Cl (Ed.6) C07C, C07D

Search Examiner  
MR R HONEYWOOD

Date of completion of Search  
14 NOVEMBER 1994

**Databases (see below)**

(i) UK Patent Office collections of GB, EP, WO and US patent specifications.

(ii) ONLINE DATABASE - CAS

Documents considered relevant following a search in respect of Claims :-  
1-18

**Categories of documents**

- |  |   |
|--|---|
| <p><b>X:</b> Document indicating lack of novelty or of inventive step.</p> <p><b>Y:</b> Document indicating lack of inventive step if combined with one or more other documents of the same category.</p> <p><b>A:</b> Document indicating technological background and/or state of the art.</p> | <p><b>P:</b> Document published on or after the declared priority date but before the filing date of the present application.</p> <p><b>E:</b> Patent document published on or after, but with priority date earlier than, the filing date of the present application.</p> <p><b>&amp;:</b> Member of the same patent family; corresponding document.</p> |
|--|---|

Category	Identity of document and relevant passages	Relevant to claim(s)
	NONE	

**Databases:** The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).